

# PS2 Antibody

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP51891

## Product Information

---

<b>Application</b>	WB
<b>Primary Accession</b>	<a href="#">P04155</a>
<b>Reactivity</b>	Human, Mouse, Rat
<b>Host</b>	Rabbit
<b>Clonality</b>	Polyclonal
<b>Calculated MW</b>	9150

## Additional Information

---

<b>Gene ID</b>	7031
<b>Other Names</b>	Trefoil factor 1, Breast cancer estrogen-inducible protein, PNR-2, Polypeptide P1A, hP1A, Protein pS2, TFF1, BCEI, PS2
<b>Dilution</b>	WB~~1:1000
<b>Format</b>	0.01M PBS, pH 7.2, 0.09% (W/V) Sodium azide, Glycerol 50%
<b>Storage</b>	Store at -20 °C.Stable for 12 months from date of receipt

## Protein Information

---

<b>Name</b>	TFF1
<b>Synonyms</b>	BCEI, PS2
<b>Function</b>	Stabilizer of the mucous gel overlying the gastrointestinal mucosa that provides a physical barrier against various noxious agents. May inhibit the growth of calcium oxalate crystals in urine.
<b>Cellular Location</b>	Secreted
<b>Tissue Location</b>	Found in stomach, with highest levels in the upper gastric mucosal cells (at protein level). Detected in goblet cells of the small and large intestine and rectum, small submucosal glands in the esophagus, mucous acini of the sublingual gland, submucosal glands of the trachea, and epithelial cells lining the exocrine pancreatic ducts but not in the remainder of the pancreas (at protein level) Scattered expression is detected in the epithelial cells of the gallbladder and submucosal glands of the vagina, and weak expression is observed in the bronchial goblet cells of the pseudostratified epithelia in the respiratory system (at protein level). Detected in urine (at protein level). Strongly expressed in breast cancer but at low levels in normal mammary

tissue. It is regulated by estrogen in MCF-7 cells. Strong expression found in normal gastric mucosa and in the regenerative tissues surrounding ulcerous lesions of gastrointestinal tract, but lower expression found in gastric cancer (at protein level).

## Background

---

Stabilizer of the mucous gel overlying the gastrointestinal mucosa that provides a physical barrier against various noxious agents. May inhibit the growth of calcium oxalate crystals in urine.

## References

---

- Jakowlew S.B.,et al.Nucleic Acids Res. 12:2861-2878(1984).  
Prud'Homme J.-F.,et al.DNA 4:11-21(1985).  
Jeltsch J.-M.,et al.Nucleic Acids Res. 15:1401-1414(1987).  
Takahashi H.,et al.FEBS Lett. 261:283-286(1990).  
Mori K.,et al.J. Biochem. 107:73-76(1990).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.