

GPR50 Antibody

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP51961

Product Information

Application	WB
Primary Accession	Q13585
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	67369

Additional Information

Gene ID	9248
Other Names	Melatonin-related receptor, G protein-coupled receptor 50, H9, GPR50
Dilution	WB~~1:1000
Format	0.01M PBS, pH 7.2, 0.09% (W/V) Sodium azide, Glycerol 50%
Storage	Store at -20 °C.Stable for 12 months from date of receipt

Protein Information

Name	GPR50
Function	G protein-coupled receptor that plays a role in numerous physiological processes including regulation of energy metabolism, neurite outgrowth or cell migration (PubMed: 19699797). Promotes self- renewal and neuronal differentiation of neural progenitor cells through activation of the NOTCH and WNT/beta-catenin signaling pathways (By similarity). Modulates the KAT5-dependent glucocorticoid receptor signaling by modulating KAT5 subcellular compartmentalisation (PubMed: 21858214). Also plays a role in the activation TGFBR1 in the absence of TGFBR2 by interfering with FKBP1A binding to TGFBR1, leading to induction of both canonical and non-canonical SMAD signaling pathways resulting in inhibition of proliferation or promotion of migration (PubMed: 29572483).
Cellular Location	Cell membrane; Multi-pass membrane protein. Postsynaptic density
Tissue Location	Hypothalamus and pituitary.

Background

Does not bind melatonin.

References

Reppert S.M.,et al.FEBS Lett. 386:219-224(1996).

Kaighin V.A.,et al.Submitted (DEC-2007) to the EMBL/GenBank/DDBJ databases.

Levoye A.,et al.EMBO J. 25:3012-3023(2006).

Bhattacharyya S.,et al.J. Lipid Res. 47:761-766(2006).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.