10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

Anti-Acid Sphingomyelinase Antibody

Catalog # AP51974

Product Information

Application	WB
Primary Accession	P17405
Reactivity	Human, Mouse, Rat, Bovine, Sheep
Host	Rabbit
Clonality	Polyclonal
Calculated MW	69936

Additional Information

Gene ID	6609
Other Names	ASM; Sphingomyelin phosphodiesterase; Acid sphingomyelinase; aSMase
Target/Specificity	KLH-conjugated synthetic peptide encompassing a sequence within the center region of human Acid Sphingomyelinase. The exact sequence is proprietary.
Dilution	WB~~ 1:1000
Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.
Storage	Store at -20 °C.Stable for 12 months from date of receipt

Protein Information

Name	SMPD1 (HGNC:11120)
Function	Converts sphingomyelin to ceramide (PubMed: 12563314, PubMed:1840600, PubMed:18815062, PubMed:25339683, PubMed:25920558, PubMed:27659707, PubMed:33163980). Exists as two enzymatic forms that arise from alternative trafficking of a single protein precursor, one that is targeted to the endolysosomal compartment, whereas the other is released extracellularly (PubMed:20807762, PubMed:21098024, PubMed:9660788). However, in response to various forms of stress, lysosomal exocytosis may represent a major source of the secretory form (PubMed:12563314, PubMed:20530211, PubMed:20807762, PubMed:22573858, PubMed:9393854).
Cellular Location	Lysosome. Lipid droplet. Secreted. Note=The secreted form is induced in a time- and dose-dependent by IL1B and TNF as well as stress and viral infection. This increase of the secreted form seems to be due to exocytosis of the lysosomal form and is Ca(2+)-dependent (PubMed:20530211, PubMed:20807762, PubMed:22573858). Secretion is dependent of

phosphorylation at Ser-510 (PubMed:17303575). Secretion is induced by inflammatory mediators such as IL1B, IFNG or TNF as well as infection with bacteria and viruses (PubMed:12563314, PubMed:20807762)

References

Schuchman E.H.,et al.J. Biol. Chem. 266:8531-8539(1991).
Newrzella D.,et al.Biol. Chem. Hoppe-Seyler 373:1233-1238(1992).
Schuchman E.H.,et al.Genomics 12:197-205(1992).
Ida H.,et al.J. Biochem. 114:15-20(1993).
Ota T.,et al.Nat. Genet. 36:40-45(2004).

Images



Western blot analysis of Acid Sphingomyelinase expression in HL60 (A) whole cell lysates. (Predicted band size: 63; 64; 69 kD; Observed band size: 70 kD)

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.