

# HUR Antibody

Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP52000

## **Product Information**

Application	WB
Primary Accession	<u>Q15717</u>
Reactivity	Human
Host	Rabbit
Clonality	Polyclona
Calculated MW	36092

## **Additional Information**

Gene ID	1994
Other Names	ELAV-like protein 1, Hu-antigen R, HuR, ELAVL1, HUR
Target/Specificity	KLH-conjugated synthetic peptide encompassing a sequence within the center region of human HUR. The exact sequence is proprietary.
Dilution	WB~~ 1:1000
Format	0.01M PBS, pH 7.2, 0.09% (W/V) Sodium azide, Glycerol 50%
Storage	Store at -20 °C.Stable for 12 months from date of receipt

### **Protein Information**

Name	ELAVL1
Synonyms	HUR
Function	RNA-binding protein that binds to the 3'-UTR region of mRNAs and increases their stability (PubMed: <u>14517288</u> , PubMed: <u>18285462</u> , PubMed: <u>31358969</u> ). Involved in embryonic stem cell (ESC) differentiation: preferentially binds mRNAs that are not methylated by N6-methyladenosine (m6A), stabilizing them, promoting ESC differentiation (By similarity). Has also been shown to be capable of binding to m6A-containing mRNAs and contributes to MYC stability by binding to m6A-containing MYC mRNAs (PubMed: <u>32245947</u> ). Binds to poly-U elements and AU-rich elements (AREs) in the 3'-UTR of target mRNAs (PubMed: <u>14731398</u> , PubMed: <u>17632515</u> , PubMed: <u>18285462</u> , PubMed: <u>23519412</u> , PubMed: <u>8626503</u> ). Binds avidly to the AU-rich element in FOS and IL3/interleukin-3 mRNAs. In the case of the FOS AU-rich element, binds to a core element of 27 nucleotides that contain AUUUA, AUUUUA, and AUUUUUA motifs. Binds preferentially to the 5'-UUUU[AG]UUU-3' motif in vitro (PubMed: <u>8626503</u> ). With ZNF385A, binds the 3'-UTR of p53/TP53 mRNA

	to control their nuclear export induced by CDKN2A. Hence, may regulate p53/TP53 expression and mediate in part the CDKN2A anti-proliferative activity. May also bind with ZNF385A the CCNB1 mRNA (By similarity). Increases the stability of the leptin mRNA harboring an AU-rich element (ARE) in its 3' UTR (PubMed: <u>29180010</u> ).
Cellular Location	Cytoplasm. Nucleus. Cytoplasm, Stress granule {ECO:0000250 UniProtKB:P70372}. Cytoplasm, P-body. Note=Translocates into the cytoplasm following phosphorylation by MAPKAPK2 (PubMed:14517288). Likewise, phosphorylation by PRKCD promotes translocation from the nucleus into the cytoplasm, where it is associated with free and cytoskeleton-bound polysomes (PubMed:18285462). Localizes to the stress granules in the presence of PLEKHN1 (By similarity). {ECO:0000250 UniProtKB:P70372, ECO:0000269 PubMed:14517288, ECO:0000269 PubMed:18285462}
Tissue Location	Ubiquitous. Detected in brain, liver, thymus and muscle.

## Background

Binds avidly to the AU-rich element in FOS and IL3/interleukin-3 mRNAs. In the case of the FOS AU-rich element, HUR binds to a core element of 27 nucleotides that contain AUUUA, AUUUUA, and AUUUUUA motifs. Binds preferentially to the 5'- UUUU[AG]UUU-3' motif in vitro. With ZNF385A, binds the 3'-UTR of p53/TP53 mRNA to control their nuclear export induced by CDKN2A. Hence, may regulate p53/TP53 expression and mediate in part the CDKN2A anti-proliferative activity. May also bind with ZNF385A the CCNB1 mRNA.

## References

Ma W.-J.,et al.J. Biol. Chem. 271:8144-8151(1996). Kalnine N.,et al.Submitted (AUG-2003) to the EMBL/GenBank/DDBJ databases. Mural R.J.,et al.Submitted (SEP-2005) to the EMBL/GenBank/DDBJ databases. Gallouzi I.-E.,et al.Science 294:1895-1901(2001). Li H.,et al.J. Biol. Chem. 277:44623-44630(2002).

#### Images



Anti-HUR Antibody at 1:1000 dilution + Jurkat whole cell lysates Lysates/proteins at 20 μg per lane. Secondary Goat Anti-Rabbit IgG, (H+L),Peroxidase conjugated at 1/10000 dilution Predicted band size : 36 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.