10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

Anti-SOX11 Antibody

Catalog # AP52016

Product Information

Application	WB, IF/IC
Primary Accession	P35716
Reactivity	Human, Mouse, Rat, Zebrafish, Chicken
Host	Rabbit
Clonality	Polyclonal
Calculated MW	46679

Additional Information

Gene ID	6664
Other Names	Transcription factor SOX-11
Target/Specificity	KLH-conjugated synthetic peptide encompassing a sequence within the N-term region of human SOX11. The exact sequence is proprietary.
Dilution	WB~~WB (1/500 - 1/1000) IF/IC~~N/A
Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.
Storage	Store at -20 °C.Stable for 12 months from date of receipt


Protein Information

Name	SOX11
Function	Transcription factor that acts as a transcriptional activator (PubMed:24886874, PubMed:26543203). Binds cooperatively with POU3F2/BRN2 or POU3F1/OCT6 to gene promoters, which enhances transcriptional activation (By similarity). Acts as a transcriptional activator of TEAD2 by binding to its gene promoter and first intron (By similarity). Plays a redundant role with SOX4 and SOX12 in cell survival of developing tissues such as the neural tube, branchial arches and somites, thereby contributing to organogenesis (By similarity).
Cellular Location	Nucleus {ECO:0000255|PROSITE-ProRule:PRU00267, ECO:0000269|PubMed:24886874, ECO:0000269|PubMed:35938035}
Tissue Location	Expressed primarily in the brain and heart, with low expression in the kidney, pancreas and muscle


References

Jay P.,et al.Genomics 29:541-545(1995).
Azuma T.,et al.DNA Res. 6:357-360(1999).
Hillier L.W.,et al.Nature 434:724-731(2005).
Goze C.,et al.Nucleic Acids Res. 21:2943-2943(1993).
Dephoure N.,et al.Proc. Natl. Acad. Sci. U.S.A. 105:10762-10767(2008).

Images



Western blot analysis of SOX11 expression in Hela (A), DLD (B) whole cell lysates. (Predicted band size: 46 kD; Observed band size: 46; 70 kD)



Immunofluorescent analysis of SOX11 staining in Hela cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a hidified chamber. Cells were washed with PBST and incubated with a AF488-conjugated secondary antibody (green) in PBS at room temperature in the dark. Phalloidin - AF594 was used to stain Actin filaments (red). DAPI was used to stain the cell nuclei (blue).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.