

CD154 Antibody

Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP52025

Product Information

Application WB Primary Accession P29965

Reactivity Human, Mouse, Rat

HostRabbitClonalityPolyclonalCalculated MW29274

Additional Information

Gene ID 959

Other Names CD40 ligand, CD40-L, T-cell antigen Gp39, TNF-related activation protein,

TRAP, Tumor necrosis factor ligand superfamily member 5, CD154, CD40 ligand, membrane form, CD40 ligand, soluble form, CD40LG, CD40L, TNFSF5,

TRAP

Dilution WB~~1:1000

Format 0.01M PBS, pH 7.2, 0.09% (W/V) Sodium azide, Glycerol 50%

Storage Store at -20 °C.Stable for 12 months from date of receipt

Protein Information

Name CD40LG

Synonyms CD40L, TNFSF5, TRAP

Function Cytokine that acts as a ligand to CD40/TNFRSF5 (PubMed: 1280226,

PubMed:31331973). Costimulates T-cell proliferation and cytokine production (PubMed:8617933). Its cross-linking on T-cells generates a costimulatory signal which enhances the production of IL4 and IL10 in conjunction with the TCR/CD3 ligation and CD28 costimulation (PubMed:8617933). Induces the activation of NF-kappa-B (PubMed:15067037, PubMed:31331973). Induces the activation of kinases MAPK8 and PAK2 in T-cells (PubMed:15067037). Induces tyrosine phosphorylation of isoform 3 of CD28 (PubMed:15067037). Mediates B-cell proliferation in the absence of co-stimulus as well as IgE production in the presence of IL4 (By similarity). Involved in immunoglobulin class switching

(By similarity).

Cellular Location Cell membrane; Single-pass type II membrane protein. Cell surface

Background

Mediates B-cell proliferation in the absence of co-stimulus as well as IgE production in the presence of IL-4. Involved in immunoglobulin class switching.

References

Graf D.,et al.Eur. J. Immunol. 22:3191-3194(1992). Hollenbaugh D.,et al.EMBO J. 11:4313-4321(1992). Aruffo A.,et al.Cell 72:291-300(1993). Spriggs M.K.,et al.J. Exp. Med. 176:1543-1550(1992). Gauchat J.-F.,et al.FEBS Lett. 315:259-266(1993).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.