

Angiopoietin 2 Rabbit pAb

Angiopoietin 2 Rabbit pAb

Catalog # AP52037

Product Information

Application	WB, IHC-P, IHC-F, IF
Primary Accession	O15123
Reactivity	Human
Predicted	Rat, Pig, Dog, Mouse, Chicken
Host	Rabbit
Clonality	Polyclonal
Calculated MW	56919
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human Angiopoietin 2
Epitope Specificity	401-496/496
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SUBCELLULAR LOCATION	Secreted.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	The protein encoded by this gene is an antagonist of angiopoietin 1 (ANGPT1) and endothelial TEK tyrosine kinase (TIE-2, TEK). The encoded protein disrupts the vascular remodeling ability of ANGPT1 and may induce endothelial cell apoptosis. Three transcript variants encoding three different isoforms have been found for this gene. [provided by RefSeq, Jul 2008]

Additional Information

Gene ID	285
Other Names	Angiopoietin-2, ANG-2, ANGPT2
Dilution	WB=1:500-2000,IHC-P=1:100-500,IHC-F=1:100-500,IF=1:100-500
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Protein Information

Name	ANGPT2
Function	Binds to TEK/TIE2, competing for the ANGPT1 binding site, and modulating

ANGPT1 signaling (PubMed:[15284220](#), PubMed:[19116766](#), PubMed:[19223473](#), PubMed:[9204896](#)). Can induce tyrosine phosphorylation of TEK/TIE2 in the absence of ANGPT1 (PubMed:[15284220](#), PubMed:[19116766](#), PubMed:[19223473](#), PubMed:[9204896](#)). In the absence of angiogenic inducers, such as VEGF, ANGPT2-mediated loosening of cell-matrix contacts may induce endothelial cell apoptosis with consequent vascular regression. In concert with VEGF, it may facilitate endothelial cell migration and proliferation, thus serving as a permissive angiogenic signal (PubMed:[15284220](#), PubMed:[19116766](#), PubMed:[19223473](#), PubMed:[9204896](#)). Involved in the regulation of lymphangiogenesis (PubMed:[32908006](#)).

Cellular Location Secreted.


Background

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.


References

Maisonpierre P.C.,et al.Science 277:55-60(1997).
 Tanaka S.,et al.J. Clin. Invest. 103:341-345(1999).
 Kim I.,et al.J. Biol. Chem. 275:18550-18556(2000).
 Ota T.,et al.Nat. Genet. 36:40-45(2004).
 Nusbaum C.,et al.Nature 439:331-335(2006).

Images



Sample:
 HepG2(Human) Cell Lysate at 30 ug
 Primary: Anti-Angiopoietin 2 (AP52037) at 1/300 dilution
 Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution
 Predicted band size: 53 kD
 Observed band size: 53 kD



Tissue/cell: human colon cancer; 4%
 Paraformaldehyde-fixed and paraffin-embedded;
 Antigen retrieval: citrate buffer (0.01M, pH 6.0), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum,C-0005) at 37°C for 20 min;
 Incubation: Anti-Ang2 Polyclonal Antibody, Unconjugated(AP52037) 1:500, overnight at 4°C, followed by conjugation to the secondary antibody(SP-0023) and DAB(C-0010) staining