

C5b-9 Rabbit pAb

C5b-9 Rabbit pAb
Catalog # AP52073

Product Information

Application	WB, IHC-P, IHC-F, IF
Primary Accession	P01031
Reactivity	Rat, Human
Predicted	Mouse
Host	Rabbit
Clonality	Polyclonal
Calculated MW	188305
Physical State	Liquid
Immunogen	Purified Human SC5b-9 Complex
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SUBCELLULAR LOCATION	Secreted.
DISEASE	Defects in C5 are the cause of complement component 5 deficiency (C5D) [MIM:609536]. A rare defect of the complement classical pathway associated with susceptibility to severe recurrent infections, predominantly by <i>Neisseria gonorrhoeae</i> or <i>Neisseria meningitidis</i> . Note=An association study of C5 haplotypes and genotypes in individuals with chronic hepatitis C virus infection shows that individuals homozygous for the C5_1 haplotype have a significantly higher stage of liver fibrosis than individuals carrying at least 1 other allele (PubMed:15995705).
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	Activation of the complement system plays a key role in normal inflammatory response to injury but may cause substantial injury when activated inappropriately. The complement system is activated either through the classical (antibody induced) or the alternative (microbial surface, polysaccharide induced) pathway, both leading to the formation of the C5b9 complex. Fluid phase binding of the multifunctional glycoprotein S protein (vitronectin) to C5b9 leads to the formation of a cytolytically inactive complex, SC5b9, which is unable to attach to cells.

Additional Information

Gene ID	727
Other Names	Complement C5, C3 and PZP-like alpha-2-macroglobulin domain-containing protein 4, Complement C5 beta chain, Complement C5 alpha chain, C5a anaphylatoxin, Complement C5b, Complement C5 alpha' chain, C5 {ECO:0000303 PubMed:1984448, ECO:0000312 HGNC:HGNC:1331}

Dilution	WB=1:500-2000,IHC-P=1:100-500,IHC-F=1:100-500,IF=1:100-500
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Protein Information

Name	C5 {ECO:0000303 PubMed:1984448, ECO:0000312 HGNC:HGNC:1331}
Function	Precursor of the C5a anaphylatoxin and complement C5b components of the complement pathways, which consist in a cascade of proteins that leads to phagocytosis and breakdown of pathogens and signaling that strengthens the adaptive immune system (PubMed: 12878586 , PubMed: 18204047 , PubMed: 30643019 , PubMed: 6554279). Activated downstream of classical, alternative, lectin and GZMK complement pathways (PubMed: 12878586 , PubMed: 18204047 , PubMed: 30643019 , PubMed: 39914456 , PubMed: 39814882 , PubMed: 6554279).
Cellular Location	Secreted. [C5a anaphylatoxin]: Secreted


Background

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

References


Stanley K.K.,et al.EMBO J. 4:375-382(1985).
 Discipio R.G.,et al.Proc. Natl. Acad. Sci. U.S.A. 81:7298-7302(1984).
 Marazziti D.,et al.Biochemistry 27:6529-6534(1988).
 Witzel-Schloemp K.,et al.Immunogenetics 48:144-147(1998).
 DiScipio R.G.,et al.J. Biol. Chem. 260:14802-14809(1985).

Images



Sample:
 Lane 1: A549 (Human) Cell Lysate at 30 ug
 Lane 2: Hela (Human) Cell Lysate at 30 ug
 Primary: Anti-C5b (AP52073) at 1/1000 dilution
 Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution
 Predicted band size: 72 kD
 Observed band size: 72 kD

Paraformaldehyde-fixed, paraffin embedded (Rat liver);
 Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer



(normal goat serum) at 37°C for 30min; Antibody incubation with (C5b-9) Polyclonal Antibody, Unconjugated (AP52073) at 1:500 overnight at 4°C, followed by a conjugated secondary (sp-0023) for 20 minutes and DAB staining.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.