

TNF alpha Rabbit pAb

TNF alpha Rabbit pAb

Catalog # AP52092

Product Information

Primary Accession	P01375
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	25644
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human TNF-alpha
Epitope Specificity	181-235/235
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SUBCELLULAR LOCATION	Cell membrane; Single-pass type II membrane protein. Tumor necrosis factor, membrane form: Membrane; Single-pass type II membrane protein. Tumor necrosis factor, soluble form: Secreted. C-domain 1: Secreted. C-domain 2: Secreted.
DISEASE	Genetic variations in TNF are a cause of susceptibility psoriatic arthritis (PSORAS) [MIM:607507]. PSORAS is an inflammatory, seronegative arthritis associated with psoriasis. It is a heterogeneous disorder ranging from a mild, non-destructive disease to a severe, progressive, erosive arthropathy. Five types of psoriatic arthritis have been defined: asymmetrical oligoarthritis characterized by primary involvement of the small joints of the fingers or toes; asymmetrical arthritis which involves the joints of the extremities; symmetrical polyarthritis characterized by a rheumatoidlike pattern that can involve hands, wrists, ankles, and feet; arthritis mutilans, which is a rare but deforming and destructive condition; arthritis of the sacroiliac joints and spine (psoriatic spondylitis).
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	This gene encodes a multifunctional proinflammatory cytokine that belongs to the tumor necrosis factor (TNF) superfamily. This cytokine is mainly secreted by macrophages. It can bind to, and thus functions through its receptors TNFRSF1A/TNFR1 and TNFRSF1B/TNFR. This cytokine is involved in the regulation of a wide spectrum of biological processes including cell proliferation, differentiation, apoptosis, lipid metabolism, and coagulation. This cytokine has been implicated in a variety of diseases, including autoimmune diseases, insulin resistance, psoriasis, rheumatoid arthritis ankylosing spondylitis, tuberculosis, autosomal dominant polycystic kidney disease, and cancer. Mutations in this gene affect susceptibility to cerebral malaria, septic shock, and Alzheimer disease. Knockout studies in mice also suggested the neuroprotective function of this cytokine. [provided by RefSeq, Aug 2020]

Additional Information

Gene ID	7124
Other Names	Tumor necrosis factor, Cachectin, TNF-alpha, Tumor necrosis factor ligand superfamily member 2, TNF-a, Tumor necrosis factor, membrane form, N-terminal fragment, NTF, Intracellular domain 1, ICD1, Intracellular domain 2, ICD2, C-domain 1, C-domain 2, Tumor necrosis factor, soluble form, TNF, TNFA, TNFSF2
Dilution	Flow-Cyt=1:50-100,ICC/IF=1:100-500
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Protein Information

Name	TNF
Synonyms	TNFA, TNFSF2
Function	Cytokine that binds to TNFRSF1A/TNFR1 and TNFRSF1B/TNFR. It is mainly secreted by macrophages and can induce cell death of certain tumor cell lines. It is potent pyrogen causing fever by direct action or by stimulation of interleukin-1 secretion and is implicated in the induction of cachexia, Under certain conditions it can stimulate cell proliferation and induce cell differentiation. Impairs regulatory T- cells (Treg) function in individuals with rheumatoid arthritis via FOXP3 dephosphorylation. Up-regulates the expression of protein phosphatase 1 (PP1), which dephosphorylates the key 'Ser-418' residue of FOXP3, thereby inactivating FOXP3 and rendering Treg cells functionally defective (PubMed: 23396208). Key mediator of cell death in the anticancer action of BCG-stimulated neutrophils in combination with DIABLO/SMAC mimetic in the RT4v6 bladder cancer cell line (PubMed: 16829952 , PubMed: 22517918 , PubMed: 23396208). Induces insulin resistance in adipocytes via inhibition of insulin-induced IRS1 tyrosine phosphorylation and insulin-induced glucose uptake. Induces GKAP42 protein degradation in adipocytes which is partially responsible for TNF-induced insulin resistance (By similarity). Plays a role in angiogenesis by inducing VEGF production synergistically with IL1B and IL6 (PubMed: 12794819). Promotes osteoclastogenesis and therefore mediates bone resorption (By similarity).
Cellular Location	Cell membrane; Single-pass type II membrane protein [Tumor necrosis factor, soluble form]; Secreted [C-domain 2]; Secreted.


Background

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.


References

Shirai T.,et al.DNA 7:193-201(1988).
Pennica D.,et al.Proc. Natl. Acad. Sci. U.S.A. 82:6060-6064(1985).
Caput D.,et al.Proc. Natl. Acad. Sci. U.S.A. 83:1670-1674(1986).
Fransen L.,et al.Nucleic Acids Res. 13:4417-4429(1985).

Images



Blank control: THP-1. Primary Antibody (green line): Rabbit Anti-TNF alpha antibody (AP52092) Dilution: 1ug/Test; Secondary Antibody : Goat anti-rabbit IgG-BF488 Dilution: 0.5ug/Test. The cells were treated with TPA (80 nM, overnight) and then treated with LPS (1 ug/mL, 18 hr/6 hr) and Brefeldin A (300 ng/mL, last 3 hr of stimulation). The cells were fixed with 4% PFA (10min at room temperature) and then permeabilized with 90% ice-cold methanol for 20 min at -20°C. The cells were then incubated in 5% BSA to block non-specific protein-protein interactions for 30 min at room temperature. Cells stained with Primary Antibody for 30 min at room temperature. The secondary antibody used for 40 min at room temperature. Acquisition of 20,000 events was performed.



LPS(100 ng/ml, 16 h) and Brefeldin A treated (5 µg/ml, 4 h) THP-1 cells; Antibody incubation with (TNF alpha) polyclonal Antibody, unconjugated (AP52092) 1:100, 90 min at 37°C; followed by conjugated Goat Anti-Rabbit IgG antibody (green, AP52092-BF488) at 37°C for 90 min, DAPI (blue, C02-04002) was used to stain the cell nuclei. PBS instead of the primary antibody was used as the blank control.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.