

Fibronectin/FN1 Rabbit pAb

Fibronectin/FN1 Rabbit pAb

Catalog # AP52116

Product Information

Application	IHC-P, IHC-F, IF
Primary Accession	P02751
Reactivity	Rat, Human, Mouse
Predicted	Dog
Host	Rabbit
Clonality	Polyclonal
Calculated MW	272320
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human Fibronectin/Ugl-Y3
Epitope Specificity	1201-1300/2386
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SUBCELLULAR LOCATION	Secreted, extracellular space, extracellular matrix.
DISEASE	Defects in FN1 are the cause of glomerulopathy with fibronectin deposits type 2 (GFND2) [MIM:601894]; also known as familial glomerular nephritis with fibronectin deposits or fibronectin glomerulopathy. GFND is a genetically heterogeneous autosomal dominant disorder characterized clinically by proteinuria, microscopic hematuria, and hypertension that leads to end-stage renal failure in the second to fifth decade of life.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	This gene encodes fibronectin, a glycoprotein present in a soluble dimeric form in plasma, and in a dimeric or multimeric form at the cell surface and in extracellular matrix. Fibronectin is involved in cell adhesion and migration processes including embryogenesis, wound healing, blood coagulation, host defense, and metastasis. The gene has three regions subject to alternative splicing, with the potential to produce 20 different transcript variants. However, the full-length nature of some variants has not been determined. [provided by RefSeq, Jul 2008].

Additional Information

Gene ID	2335
Other Names	Fibronectin, FN, Cold-insoluble globulin, CIG, Anastellin, Ugl-Y1, Ugl-Y2, Ugl-Y3, FN1 (HGNC:3778), FN
Dilution	IHC-P=1:100-500,IHC-F=1:100-500,IF=1:200-800
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When

reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Protein Information

Name	FN1 (HGNC:3778)
Synonyms	FN
Function	Fibronectins bind cell surfaces and various compounds including collagen, fibrin, heparin, DNA, and actin (PubMed: 3024962 , PubMed: 3593230 , PubMed: 3900070 , PubMed: 7989369). Fibronectins are involved in cell adhesion, cell motility, opsonization, wound healing, and maintenance of cell shape (PubMed: 3024962 , PubMed: 3593230 , PubMed: 3900070 , PubMed: 7989369). Involved in osteoblast compaction through the fibronectin fibrillogenesis cell-mediated matrix assembly process, essential for osteoblast mineralization (By similarity). Participates in the regulation of type I collagen deposition by osteoblasts (By similarity). Acts as a ligand for the LILRB4 receptor, inhibiting FCGR1A/CD64-mediated monocyte activation (PubMed: 34089617).
Cellular Location	Secreted, extracellular space, extracellular matrix. Secreted {ECO:0000250 UniProtKB:P11276}
Tissue Location	Expressed in the inner limiting membrane and around blood vessels in the retina (at protein level) (PubMed:29777959) Plasma FN (soluble dimeric form) is secreted by hepatocytes. Cellular FN (dimeric or cross-linked multimeric forms), made by fibroblasts, epithelial and other cell types, is deposited as fibrils in the extracellular matrix. Ugl-Y1, Ugl-Y2 and Ugl-Y3 are found in urine (PubMed:17614963).

Background


This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

References


Church D.M.,et al.PLoS Biol. 7:E1000112-E1000112(2009).
Mural R.J.,et al.Submitted (JUL-2005) to the EMBL/GenBank/DDBJ databases.
Polly P.,et al.Gene 137:353-354(1993).
Talts J.F.,et al.J. Cell Sci. 108:2153-2162(1995).
Gorski G.,et al.Submitted (DEC-1995) to the EMBL/GenBank/DDBJ databases.

Images

Paraformaldehyde-fixed, paraffin embedded (Rat colon);
Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Incubation: Anti-Fibronectin Polyclonal Antibody, conjugated (AP52116-AF647) 1:400, 1.5 hours at 37°C; DAPI (5ug/ml,



blue, C-0033) was used to stain the cell nuclei.



Tissue/cell: rat intestine tissue; 4%
 Paraformaldehyde-fixed and paraffin-embedded;
 Antigen retrieval: citrate buffer (0.01M, pH 6.0), Boiling
 bathing for 15min; Block endogenous peroxidase by 3%
 Hydrogen peroxide for 30min; Blocking buffer (normal
 goat serum,C-0005) at 37°C for 20 min;
 Incubation: Anti-Fibronectin/FN1 Polyclonal Antibody,
 Unconjugated(AP52116) 1:500, overnight at 4°C, followed
 by conjugation to the secondary antibody(SP-0023) and
 DAB(C-0010) staining

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.