

IGF2R/M6PR Rabbit pAb

IGF2R/M6PR Rabbit pAb

Catalog # AP52118

Product Information

Application	IHC-P, IHC-F, IF
Primary Accession	P11717
Reactivity	Human, Mouse
Predicted	Rat, Pig, Dog, Rabbit, Chicken, Horse
Host	Rabbit
Clonality	Polyclonal
Calculated MW	274375
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human IGF2R
Epitope Specificity	2201-2400/2491
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SUBCELLULAR LOCATION	Lysosome membrane. Colocalized with DPP4 in internalized cytoplasmic vesicles adjacent to the cell surface.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	Transport of phosphorylated lysosomal enzymes from the Golgi complex and the cell surface to lysosomes. Lysosomal enzymes bearing phosphomannosyl residues bind specifically to mannose-6-phosphate receptors in the Golgi apparatus and the resulting receptor-ligand complex is transported to an acidic prelysosomal compartment where the low pH mediates the dissociation of the complex. This receptor also binds IGF2. Acts as a positive regulator of T-cell coactivation, by binding DPP4.

Additional Information

Gene ID	3482
Other Names	Cation-independent mannose-6-phosphate receptor, CI Man-6-P receptor, CI-MPR, M6PR, 300 kDa mannose 6-phosphate receptor, MPR 300, Insulin-like growth factor 2 receptor, Insulin-like growth factor II receptor, IGF-II receptor, M6P/IGF2 receptor, M6P/IGF2R, CD222, IGF2R, MPRI
Dilution	IHC-P=1:100-500,IHC-F=1:100-500,IF=1:100-500,Flow-Cyt=5ug/Test
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Protein Information

Name IGF2R

Synonyms MPRI

Function Mediates the transport of phosphorylated lysosomal enzymes from the Golgi complex and the cell surface to lysosomes (PubMed:[18817523](#), PubMed:[2963003](#)). Lysosomal enzymes bearing phosphomannosyl residues bind specifically to mannose-6-phosphate receptors in the Golgi apparatus and the resulting receptor-ligand complex is transported to an acidic prelysosomal compartment where the low pH mediates the dissociation of the complex (PubMed:[18817523](#), PubMed:[2963003](#)). The receptor is then recycled back to the Golgi for another round of trafficking through its binding to the retromer (PubMed:[18817523](#)). This receptor also binds IGF2 (PubMed:[18046459](#)). Acts as a positive regulator of T-cell coactivation by binding DPP4 (PubMed:[10900005](#)).

Cellular Location Golgi apparatus membrane; Single-pass type I membrane protein. Endosome membrane; Single-pass type I membrane protein. Note=Mainly localized in the Golgi at steady state and not detectable in lysosome (PubMed:[18817523](#)) Colocalized with DPP4 in internalized cytoplasmic vesicles adjacent to the cell surface (PubMed:[10900005](#)).


Background

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

References


Morgan D.O.,et al.Nature 329:301-307(1987).
Oshima A.,et al.J. Biol. Chem. 263:2553-2562(1988).
Gemma A.,et al.Submitted (NOV-1998) to the EMBL/GenBank/DDBJ databases.
Killian J.K.,et al.Mamm. Genome 10:74-77(1999).
Mungall A.J.,et al.Nature 425:805-811(2003).

Images



Blank control(blue): Raji (fixed with 2% paraformaldehyde (10 min)). Primary Antibody:Rabbit Anti-IGF2R M6PR antibody(AP52118), Dilution: 5 µg in 100 µL 1X PBS containing 0.5% BSA; Isotype Control Antibody: Rabbit IgG(orange) ,used under the same conditions); Secondary Antibody: Goat anti-rabbit IgG-PE(white blue), Dilution: 1:200 in 1 X PBS containing 0.5% BSA.

Paraformaldehyde-fixed, paraffin embedded (human meningioma); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous



peroxidase by 3% hydrogen peroxide for 20 minutes;
Blocking buffer (normal goat serum) at 37°C for 30min;
Antibody incubation with (IGF2R) Polyclonal Antibody,
Unconjugated (AP52118) at 1:400 overnight at 4°C,
followed by operating according to SP Kit(Rabbit)
(sp-0023) instructions and DAB staining.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.