

# human secretory IgA Rabbit pAb

human secretory IgA Rabbit pAb

Catalog # AP52174

## Product Information

---

|                                |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                               |
|--------------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <b>Application</b>             | WB                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                            |
| <b>Primary Accession</b>       | <a href="#">P01876</a>                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                        |
| <b>Reactivity</b>              | Human                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                         |
| <b>Host</b>                    | Rabbit                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                        |
| <b>Clonality</b>               | Polyclonal                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                    |
| <b>Calculated MW</b>           | 42849                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                         |
| <b>Physical State</b>          | Liquid                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                        |
| <b>Immunogen</b>               | Purified human sIgA                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                           |
| <b>Isotype</b>                 | IgG                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                           |
| <b>Purity</b>                  | affinity purified by Protein A                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                |
| <b>Buffer</b>                  | 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                             |
| <b>SUBCELLULAR LOCATION</b>    | Secreted.                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                     |
| <b>Important Note</b>          | This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.                                                                                                                                                                                                                                                                                                                                                                                                                                                                     |
| <b>Background Descriptions</b> | Monomeric IgA constitutes 5-15 % of the serum immunoglobulins whereas dimeric IgA is localized to mucosa surfaces such as saliva, gastrointestinal secretion, bronchial fluids and milk. Mucosal IgA plays a major role in host defence by neutralising infectious agents at mucosal surfaces. The production is usually local and antigen specific IgA producing B-cells can be found in regions under the lamina propria where they mature into dimeric IgA producing plasma cells. IgA deficiency is the most common immunodeficiency that may affect both serum and mucosal produced IgA. |

## Additional Information

---

|                    |                                                                                                                                                                                                                                                   |
|--------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <b>Other Names</b> | Immunoglobulin heavy constant alpha 1 {ECO:0000303   PubMed:11340299, ECO:0000303   Ref.13}, Ig alpha-1 chain C region, Ig alpha-1 chain C region BUR, Ig alpha-1 chain C region TRO, IGHA1 {ECO:0000303   PubMed:11340299, ECO:0000303   Ref.13} |
| <b>Dilution</b>    | WB=1:500-2000                                                                                                                                                                                                                                     |
| <b>Storage</b>     | Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.                                                   |

## Protein Information

---

|             |                                                             |
|-------------|-------------------------------------------------------------|
| <b>Name</b> | IGHA1 {ECO:0000303   PubMed:11340299, ECO:0000303   Ref.13} |
|-------------|-------------------------------------------------------------|

|                          |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                               |
|--------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <b>Function</b>          | Constant region of immunoglobulin heavy chains. Immunoglobulins, also known as antibodies, are membrane-bound or secreted glycoproteins produced by B lymphocytes. In the recognition phase of humoral immunity, the membrane-bound immunoglobulins serve as receptors which, upon binding of a specific antigen, trigger the clonal expansion and differentiation of B lymphocytes into immunoglobulins-secreting plasma cells. Secreted immunoglobulins mediate the effector phase of humoral immunity, which results in the elimination of bound antigens (PubMed: <a href="#">20176268</a> , PubMed: <a href="#">22158414</a> ). The antigen binding site is formed by the variable domain of one heavy chain, together with that of its associated light chain. Thus, each immunoglobulin has two antigen binding sites with remarkable affinity for a particular antigen. The variable domains are assembled by a process called V-(D)-J rearrangement and can then be subjected to somatic hypermutations which, after exposure to antigen and selection, allow affinity maturation for a particular antigen (PubMed: <a href="#">17576170</a> , PubMed: <a href="#">20176268</a> ). Ig alpha is the major immunoglobulin class in body secretions (PubMed: <a href="#">2241915</a> ). |
| <b>Cellular Location</b> | [Isoform 1]: Secreted                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                         |


## Background

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

## References

Uchida Y.,et al.Neuron 7:337-347(1991).  
Palmiter R.D.,et al.Proc. Natl. Acad. Sci. U.S.A. 89:6333-6337(1992).  
Tsuji S.,et al.EMBO J. 11:4843-4850(1992).  
Naruse S.,et al.Gene 144:283-287(1994).  
Amoureux M.C.,et al.Biochem. Biophys. Res. Commun. 214:639-645(1995).

## Images



### Sample:

Lane 1: U2os (Human) Cell Lysate at 30 ug

Lane 2: MCF-7 (Human) Cell Lysate at 30 ug

Primary: Anti-human secretory IgA (AP52174) at 1/1000 dilution

Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution

Observed band size: 62 kD

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.