

HBsAg Goat pAb

HBsAg Goat pAb
Catalog # AP52176

Product Information

Application	WB, IHC-P, IHC-F, IF
Host	Goat
Clonality	Polyclonal
Physical State	Liquid
Immunogen	Recombinant HBsAg protein
Isotype	IgG
Purity	affinity purified by Protein G
Buffer	0.01M PBS (pH7.4) with 0.02% Proclin300.
SUBCELLULAR LOCATION	Virion membrane.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	人乙型肝炎表面抗原抗体

Additional Information

Other Names	Microtubule-associated protein 4; MAP-4; MAP4
Dilution	WB=1:500-2000,IHC-P=1:100-500,IHC-F=1:100-500,IF=1:100-500,ELISA=1:5000-10000
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Background


This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

References


West R.R.,et al.J. Biol. Chem. 266:21886-21896(1991).
Chapin S.J.,et al.Biochemistry 34:2289-2301(1995).
Ota T.,et al.Nat. Genet. 36:40-45(2004).
Totoki Y.,et al.Submitted (MAR-2005) to the EMBL/GenBank/DDBJ databases.
Muzny D.M.,et al.Nature 440:1194-1198(2006).

Images

Goat Anti-Human HBsAg Antibody Bind with Human HBsAg protein, His tag



Measured by its binding ability in a indirect ELISA.
Immobilized recombinant HBsAg protein (Cat. AP52176) at 2 µg/mL (100 µL/well) can bind Goat Anti-Human HBsAg Antibody, the EC50 is 29.14 ng/mL.



Sample:
Lane 1: Recombinant HBsAg protein(AP52176)
Primary: Anti-HBsAg (AP52176) at 1/1000 dilution
Secondary: IRDye800CW Donkey anti-Goat IgG at 1/20000 dilution
Predicted band size: 44 kDa
Observed band size: 20 kDa

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.