

Neurocan Rabbit pAb

Neurocan Rabbit pAb

Catalog # AP52225

Product Information

Application	IHC-P, IHC-F, IF
Primary Accession	O14594
Reactivity	Rat, Mouse
Predicted	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	143093
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human Neurocan
Epitope Specificity	1185-1257/1257
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SUBCELLULAR LOCATION	Secreted.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	Neurocan is the major soluble chondroitin sulfate proteoglycan in the brain. It is thought to play a functional role in axonal growth and guidance and in the establishment of specific neural pathways during embryonic brain development. Neurocan expression in the brain is developmentally regulated. Early on the major form of neurocan consists of a 245kD core protein with approximately two chondroitin sulfate glycosaminoglycan chains of 22kD each. Later neurocan comprises a 180kD core protein Both forms of neurocan contain only chondroitin 4-sulfate glycosaminoglycan chains. By virtue of their high expression at sites of neural damage and trauma, chondroitin sulfate proteoglycans, including neurocan, are thought to inhibit successful nerve regeneration. Alternative names: Chondroitin sulfate proteoglycan 3; Cspg3; NCAN; NEUR; Neurocan core protein.

Additional Information

Gene ID	1463
Other Names	Neurocan core protein, Chondroitin sulfate proteoglycan 3, NCAN, CSPG3, NEUR
Dilution	IHC-P=1:100-500,IHC-F=1:100-500,IF=1:100-500
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Protein Information

Name	NCAN
Synonyms	CSPG3, NEUR
Function	May modulate neuronal adhesion and neurite growth during development by binding to neural cell adhesion molecules (NG-CAM and N- CAM). Chondroitin sulfate proteoglycan; binds to hyaluronic acid.
Cellular Location	Secreted.
Tissue Location	Detected in cerebrospinal fluid (at protein level) (PubMed:25326458). Brain.


Background

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

References

Rauch U.,et al.Genomics 28:405-410(1995).
Carninci P.,et al.Science 309:1559-1563(2005).

Images



Positive sample : Paraformaldehyde-fixed, paraffin embedded (rat brain);
Negative sample : Paraformaldehyde-fixed, paraffin embedded (mouse liver);
Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (Neurocan) Polyclonal Antibody, Unconjugated (AP52225) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.