

Dopamine Transporter Rabbit pAb

Dopamine Transporter Rabbit pAb

Catalog # AP52244

Product Information

Application	WB
Primary Accession	Q01959
Reactivity	Rat
Predicted	Human, Mouse
Host	Rabbit
Clonality	Polyclonal
Calculated MW	68495
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human DAT1
Epitope Specificity	579-620/620
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SUBCELLULAR LOCATION	Membrane; Multi-pass membrane protein.
DISEASE	Defects in SLC6A3 are the cause of dystonia-parkinsonism infantile (DYTPRI) [MIM:613135]. It is a neurodegenerative disorder characterized by infantile onset of parkinsonism and dystonia. Other neurologic features include global developmental delay, bradikinesia and pyramidal tract signs.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	This gene encodes a dopamine transporter which is a member of the sodium- and chloride-dependent neurotransmitter transporter family. The 3' UTR of this gene contains a 40 bp tandem repeat, referred to as a variable number tandem repeat or VNTR, which can be present in 3 to 11 copies. Variation in the number of repeats is associated with idiopathic epilepsy, attention-deficit hyperactivity disorder, dependence on alcohol and cocaine, susceptibility to Parkinson disease and protection against nicotine dependence.[provided by RefSeq, Nov 2009]

Additional Information

Gene ID	6531
Other Names	Sodium-dependent dopamine transporter, DA transporter, DAT, Solute carrier family 6 member 3, SLC6A3, DAT1
Dilution	WB=1:500-2000
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Protein Information

Name	SLC6A3
Synonyms	DAT1
Function	Mediates sodium- and chloride-dependent transport of dopamine (PubMed: 10375632 , PubMed: 11093780 , PubMed: 1406597 , PubMed: 15505207 , PubMed: 19478460 , PubMed: 39112701 , PubMed: 39112703 , PubMed: 39112705 , PubMed: 8302271). Also mediates sodium- and chloride-dependent transport of norepinephrine (also known as noradrenaline) (By similarity). Regulator of light-dependent retinal hyaloid vessel regression, downstream of OPN5 signaling (By similarity).
Cellular Location	Cell membrane; Multi-pass membrane protein. Cell projection, neuron projection {ECO:0000250 UniProtKB:P23977}. Cell projection, axon. Note=Localizes to neurite tips in neuronal cells (By similarity). Colocalizes with SEPTIN4 at axon terminals, especially at the varicosities (By similarity). {ECO:0000250 UniProtKB:P23977, ECO:0000250 UniProtKB:Q61327}
Tissue Location	Highly expressed in substantia nigra (PubMed:7637582). Expressed in axonal varicosities in dopaminergic nerve terminals (at protein level) (PubMed:17296554). Expressed in the striatum (at protein level) (PubMed:17296554)


Background

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

References

Vandenbergh D.J.,et al.Brain Res. Mol. Brain Res. 15:161-166(1992).
Giros B.,et al.Mol. Pharmacol. 42:383-390(1992).
Pristupa Z.B.,et al.Mol. Pharmacol. 45:125-135(1994).
Kawarai T.,et al.Gene 195:11-18(1997).
Vandenbergh D.J.,et al.Mol. Psychiatry 5:283-292(2000).

Images



Sample:

Lane 1: Rat Cerebrum tissue lysates

Lane 2: Rat Liver tissue lysates(Negative control)

Primary: Anti-Dopamine Transporter (AP52244) at 1/1000 dilution

Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution

Predicted band size: 68 kDa

Observed band size: 73 kDa

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.