

Cyclooxygenase 2 Rabbit pAb

Cyclooxygenase 2 Rabbit pAb Catalog # AP52246

Product Information

Application WB, IHC-P, IHC-F, IF

Primary Accession <u>P35354</u>

Reactivity Human, Mouse

Host Rabbit
Clonality Polyclonal
Calculated MW 68996
Physical State Liquid

Immunogen KLH conjugated synthetic peptide derived from human Cyclooxygenase 2

Epitope Specificity 501-604/604

Isotype IgG

Purity affinity purified by Protein A

Buffer 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.

SUBCELLULAR LOCATION Microsome membrane; Peripheral membrane protein. Endoplasmic reticulum

membrane; Peripheral membrane protein. Post-translational modifications: S-nitrosylation by NOS2 (iNOS) activates enzme activity. S-nitrosylation may

take place on different Cys residues in addition to Cys-561.

SIMILARITY Belongs to the prostaglandin G/H synthase family. Contains 1 EGF-like

domain.

SUBUNIT Homodimer.

Post-translational S-nitrosylation by NOS2 (iNOS) activates enzme activity. S-nitrosylation may

modifications take place on different Cys residues in addition to Cys-561.

Important Note This product as supplied is intended for research use only, not for use in

human, therapeutic or diagnostic applications.

Background Descriptions Prostaglandin-endoperoxide synthase (PTGS), also known as cyclooxygenase,

is the key enzyme in prostaglandin biosynthesis, and acts both as a dioxygenase and as a peroxidase. There are two isozymes of PTGS: a

constitutive PTGS1 and an inducible PTGS2, which differ in their regulation of expression and tissue distribution. This gene encodes the inducible isozyme. It is regulated by specific stimulatory events, suggesting that it is responsible for the prostanoid biosynthesis involved in inflammation and mitogenesis.

[provided by RefSeq, Feb 2009]

Additional Information

Gene ID 5743

Other Names Prostaglandin G/H synthase 2, 1.14.99.1, Cyclooxygenase-2, COX-2, PHS II,

Prostaglandin H2 synthase 2, PGH synthase 2, PGHS-2,

Prostaglandin-endoperoxide synthase 2, PTGS2 (HGNC:9605)

Dilution WB=1:500-2000,IHC-P=1:100-500,IHC-F=1:100-500,ICC/IF=1:100-500,IF=1:100-

Storage

Store at -20 $^{\circ}$ C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 $^{\circ}$ C.

Protein Information

Name

PTGS2 (HGNC:9605)

Function

Dual cyclooxygenase and peroxidase in the biosynthesis pathway of prostanoids, a class of C20 oxylipins mainly derived from arachidonate ((5Z,8Z,11Z,14Z)-eicosatetraenoate, AA, C20:4(n-6)), with a particular role in the inflammatory response (PubMed: 11939906, PubMed: 16373578, PubMed:19540099, PubMed:22942274, PubMed:26859324, PubMed:<u>27226593</u>, PubMed:<u>7592599</u>, PubMed:<u>7947975</u>, PubMed:<u>9261177</u>). The cyclooxygenase activity oxygenates AA to the hydroperoxy endoperoxide prostaglandin G2 (PGG2), and the peroxidase activity reduces PGG2 to the hydroxy endoperoxide prostaglandin H2 (PGH2), the precursor of all 2-series prostaglandins and thromboxanes (PubMed:16373578, PubMed:22942274, PubMed:<u>26859324</u>, PubMed:<u>27226593</u>, PubMed:<u>7592599</u>, PubMed:<u>7947975</u>, PubMed: 9261177). This complex transformation is initiated by abstraction of hydrogen at carbon 13 (with S- stereochemistry), followed by insertion of molecular O2 to form the endoperoxide bridge between carbon 9 and 11 that defines prostaglandins. The insertion of a second molecule of O2 (bis-oxygenase activity) yields a hydroperoxy group in PGG2 that is then reduced to PGH2 by two electrons (PubMed:16373578, PubMed:22942274, PubMed:26859324, PubMed:27226593, PubMed:7592599, PubMed:7947975, PubMed: 9261177). Similarly catalyzes successive cyclooxygenation and peroxidation of dihomo-gamma-linoleate (DGLA, C20:3(n-6)) and eicosapentaenoate (EPA, C20:5(n-3)) to corresponding PGH1 and PGH3, the precursors of 1- and 3-series prostaglandins (PubMed: 11939906, PubMed: 19540099). In an alternative pathway of prostanoid biosynthesis, converts 2-arachidonovl lysophopholipids to prostanoid lysophopholipids, which are then hydrolyzed by intracellular phospholipases to release free prostanoids (PubMed: <u>27642067</u>). Metabolizes 2-arachidonoyl glycerol yielding the glyceryl ester of PGH2, a process that can contribute to pain response (PubMed: 22942274). Generates lipid mediators from n-3 and n-6 polyunsaturated fatty acids (PUFAs) via a lipoxygenase-type mechanism. Oxygenates PUFAs to hydroperoxy compounds and then reduces them to corresponding alcohols (PubMed: 11034610, PubMed: 11192938, PubMed: 9048568, PubMed: 9261177). Plays a role in the generation of resolution phase interaction products (resolvins) during both sterile and infectious inflammation (PubMed: 12391014). Metabolizes docosahexaenoate (DHA, C22:6(n-3)) to 17R-HDHA, a precursor of the D-series resolvins (RvDs) (PubMed: 12391014). As a component of the biosynthetic pathway of E-series resolvins (RvEs), converts eicosapentaenoate (EPA, C20:5(n-3)) primarily to 18S-HEPE that is further metabolized by ALOX5 and LTA4H to generate 18S-RvE1 and 18S-RvE2 (PubMed:21206090). In vascular endothelial cells, converts docosapentaenoate (DPA, C22:5(n-3)) to 13R- HDPA, a precursor for 13-series resolvins (RvTs) shown to activate macrophage phagocytosis during bacterial infection (PubMed: 26236990). In activated leukocytes, contributes to oxygenation of hydroxyeicosatetraenoates (HETE) to diHETES (5,15-diHETE and 5,11- diHETE) (PubMed:22068350, PubMed:26282205). Can also use linoleate (LA, (9Z,12Z)-octadecadienoate, C18:2(n-6)) as substrate and produce hydroxyoctadecadienoates (HODEs) in a regio- and stereospecific manner, being (9R)-HODE ((9R)-hydroxy-(10E,12Z)-octadecadienoate) and (13S)- HODE ((13S)-hydroxy-(9Z,11E)-octadecadienoate) its major products (By similarity).

During neuroinflammation, plays a role in neuronal secretion of specialized preresolving mediators (SPMs) 15R-lipoxin A4 that regulates phagocytic microglia (By similarity).

Cellular Location

Microsome membrane; Peripheral membrane protein. Endoplasmic reticulum membrane; Peripheral membrane protein. Nucleus inner membrane; Peripheral membrane protein. Nucleus outer membrane; Peripheral membrane protein. Note=Detected on the lumenal side of the endoplasmic reticulum and nuclear envelope


Background

Prostaglandin-endoperoxide synthase (PTGS), also known as cyclooxygenase, is the key enzyme in prostaglandin biosynthesis, and acts both as a dioxygenase and as a peroxidase. There are two isozymes of PTGS: a constitutive PTGS1 and an inducible PTGS2, which differ in their regulation of expression and tissue distribution. This gene encodes the inducible isozyme. It is regulated by specific stimulatory events, suggesting that it is responsible for the prostanoid biosynthesis involved in inflammation and mitogenesis. [provided by RefSeq, Feb 2009]


References

Kujubu D.A., et al.J. Biol. Chem. 266:12866-12872(1991). Fletcher B.S., et al.J. Biol. Chem. 267:4338-4344(1992). Ryseck R.-P., et al.Cell Growth Differ. 3:443-450(1992). O'Banion M.K., et al. Proc. Natl. Acad. Sci. U.S.A. 89:4888-4892(1992). Carninci P., et al. Science 309:1559-1563(2005).

Images



25 ug total protein per lane of various lysates (see on figure) probed with Cyclooxygenase 2 polyclonal antibody, unconjugated (AP52246) at 1:1000 dilution and 4°C overnight incubation. Followed by conjugated secondary antibody incubation at r.t. for 60 min.



Paraformaldehyde-fixed, paraffin embedded Human Esophagus Cancer; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; Antibody incubation with Cyclooxygenase 2 Polyclonal Antibody, Unconjugated (AP52246) at 1:200 overnight at 4°C, followed by conjugation to the AP52246-HRP and DAB (C-0010) staining.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.