

TRAF2 Rabbit pAb

TRAF2 Rabbit pAb
Catalog # AP52255

Product Information

Application	WB
Primary Accession	Q12933
Reactivity	Human
Predicted	Rat, Mouse
Host	Rabbit
Clonality	Polyclonal
Calculated MW	55859
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human TRAF2
Epitope Specificity	401-501/501
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SUBCELLULAR LOCATION	Cytoplasm.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	The induction of the various cellular responses mediated by tumor necrosis factor is initiated by its interaction with the TNFR1 and TNFR2 cell surface receptors. TRAF2 interacts with the cytoplasmic domain of TNFR2. TRAF2 is a common signal transducer for TNFR2 and CD40 that mediates activation of NF kappa B.

Additional Information

Gene ID	7186
Other Names	TNF receptor-associated factor 2, 2.3.2.27, E3 ubiquitin-protein ligase TRAF2, RING-type E3 ubiquitin transferase TRAF2, Tumor necrosis factor type 2 receptor-associated protein 3, TRAF2 {ECO:0000303 PubMed:28489822, ECO:0000312 HGNC:HGNC:12032}
Dilution	WB=1:500-2000,Flow-Cyt=1 µg/Test
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Protein Information

Name TRAF2 {ECO:0000303 | PubMed:28489822, ECO:0000312 | HGNC:HGNC:12032}

Function E3 ubiquitin-protein ligase that regulates activation of NF- kappa-B and JNK and plays a central role in the regulation of cell survival and apoptosis (PubMed:[10346818](#), PubMed:[11784851](#), PubMed:[12917689](#), PubMed:[15383523](#), PubMed:[18981220](#), PubMed:[19150425](#), PubMed:[19810754](#), PubMed:[19918265](#), PubMed:[19937093](#), PubMed:[20047764](#), PubMed:[20064526](#), PubMed:[20385093](#), PubMed:[20577214](#), PubMed:[22212761](#)). Catalyzes 'Lys-63'-linked ubiquitination of target proteins, such as BIRC3, IKBKE, MLST8, RIPK1 and TICAM1 (PubMed:[23453969](#), PubMed:[28489822](#)). Is an essential constituent of several E3 ubiquitin- protein ligase complexes, where it promotes the ubiquitination of target proteins by bringing them into contact with other E3 ubiquitin ligases (PubMed:[15383523](#), PubMed:[18981220](#)). Regulates BIRC2 and BIRC3 protein levels by inhibiting their autoubiquitination and subsequent degradation; this does not depend on the TRAF2 RING-type zinc finger domain (PubMed:[11907583](#), PubMed:[19506082](#)). Plays a role in mediating activation of NF-kappa-B by EIF2AK2/PKR (PubMed:[15121867](#)). In complex with BIRC2 or BIRC3, promotes ubiquitination of IKBKE (PubMed:[23453969](#)). Acts as a regulator of mTORC1 and mTORC2 assembly by mediating 'Lys-63'-linked ubiquitination of MLST8, thereby inhibiting formation of the mTORC2 complex, while facilitating assembly of the mTORC1 complex (PubMed:[28489822](#)). Required for normal antibody isotype switching from IgM to IgG (By similarity).

Cellular Location Cytoplasm


Background

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

References

Song H.Y.,et al.Biochem. J. 309:825-829(1995).
Ota T.,et al.Nat. Genet. 36:40-45(2004).
Bechtel S.,et al.BMC Genomics 8:399-399(2007).
Humphray S.J.,et al.Nature 429:369-374(2004).
Mural R.J.,et al.Submitted (JUL-2005) to the EMBL/GenBank/DDBJ databases.

Images




Blank control: U937(blue). Primary Antibody: Rabbit Anti-TRAF2 antibody(AP52255), Dilution: 1 µg in 100 µL 1X PBS containing 0.5% BSA; Isotype Control Antibody: Rabbit IgG (orange) ,used under the same conditions. Secondary Antibody: Goat anti-rabbit IgG-PE(white blue), Dilution: 1:200 in 1 X PBS containing 0.5% BSA.

Protocol

The cells were fixed with 2% paraformaldehyde (10 min).Primary antibody (AP52255, 1 µg /1x10⁶ cells) were incubated for 30 min on the ice, followed by 1 X PBS containing 0.5% BSA + 10% goat serum (15 min) to block non-specific protein-protein interactions. Then the Goat

Anti-rabbit IgG/PE antibody was added into the blocking buffer mentioned above to react with the primary antibody at 1/200 dilution for 30 min on ice. Acquisition of 20,000 events was performed.



Sample: HeLa Cell (Human) Lysate at 40 ug
Primary: Anti- TRAF2 (AP52255) at 1/300 dilution
Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution
Predicted band size: 56 kD
Observed band size: 56 kD

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.