

LAG3 Rabbit pAb

LAG3 Rabbit pAb
Catalog # AP52277

Product Information

Primary Accession	P18627
Reactivity	Mouse
Predicted	Rat, Pig, Human, Rabbit, Horse
Host	Rabbit
Clonality	Polyclonal
Calculated MW	57449
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human LAG-3
Epitope Specificity	201-300/525
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SUBCELLULAR LOCATION	Membrane; Single-pass type I membrane protein.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	Lymphocyte-activation protein 3 belongs to Ig superfamily and contains 4 extracellular Ig-like domains. The LAG3 gene contains 8 exons. The sequence data, exon/intron organization, and chromosomal localization all indicate a close relationship of LAG3 to CD4. [provided by RefSeq, Jul 2008]

Additional Information

Gene ID	3902
Other Names	Lymphocyte activation gene 3 protein, LAG-3, CD223, Secreted lymphocyte activation gene 3 protein, sLAG-3, LAG3 {ECO:0000303 PubMed:35761082, ECO:0000312 HGNC:HGNC:6476}
Dilution	Flow-Cyt=1 μ g /test
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Protein Information

Name	LAG3 {ECO:0000303 PubMed:35761082, ECO:0000312 HGNC:HGNC:6476}
Function	[Lymphocyte activation gene 3 protein]: Inhibitory receptor on antigen

activated T-cells (PubMed:[20421648](#), PubMed:[35761082](#), PubMed:[7805750](#), PubMed:[8647185](#)). Delivers inhibitory signals upon binding to ligands, such as MHC class II, its main ligand present at the surface of antigen-presenting cells (APCs), and FGL1, which is secreted by hepatocytes and certain types of tumor cells (PubMed:[30580966](#), PubMed:[32920841](#), PubMed:[35761082](#), PubMed:[39671469](#), PubMed:[7589152](#), PubMed:[8647185](#), PubMed:[9159144](#)). Ligand-binding initiates a signaling that inhibits the T-cell receptor (TCR) in the immunological synapse, preventing T-cell activation (PubMed:[40101708](#)). Mechanistically, ligand-binding promotes (1) ubiquitination of the KIEELE motif, unleashing the RRFSALE motif from the membrane and (2) leading to the formation of condensates with the TCR component CD3E, thereby disrupting the association between CD3E and LCK and preventing TCR activation (PubMed:[40101708](#), PubMed:[40592325](#)). May inhibit antigen-specific T-cell activation in synergy with PDCD1/PD-1 (By similarity). Negatively regulates the proliferation, activation, effector function and homeostasis of both CD8(+) and CD4(+) T-cells (PubMed:[20421648](#), PubMed:[7805750](#), PubMed:[8647185](#)). Also mediates immune tolerance: constitutively expressed on a subset of regulatory T-cells (Tregs) and contributes to their suppressive function (By similarity). Also acts as a negative regulator of plasmacytoid dendritic cell (pDCs) activation (By similarity).

Cellular Location

[Lymphocyte activation gene 3 protein]: Cell membrane; Single-pass type I membrane protein. Note=Clusters on the T-cell surface following ligand-binding

Tissue Location

Primarily expressed in activated T-cells and a subset of natural killer (NK) cells.


Background

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

References

Haudebourg T.,et al.Submitted (MAR-2006) to the EMBL/GenBank/DDBJ databases.

Images



Blank control:CtII-2.

Primary Antibody (green line): Rabbit Anti-LAG-3 antibody (AP52277)

Dilution: 1ug/Test;

Secondary Antibody : Goat anti-rabbit IgG-FITC

Dilution: 0.5ug/Test.

Protocol

The cells were incubated in 5%BSA to block non-specific protein-protein interactions for 30 min at room temperature .Cells stained with Primary Antibody for 30 min at room temperature. The secondary antibody used for 40 min at room temperature. Acquisition of 20,000 events was performed.