10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

# GMNN Antibody (Center Y111)

Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP5227c

## Product Information

| | |
|---|---|
| **Application** | WB, IHC-P, E |
| **Primary Accession** | O75496 |
| **Other Accession** | O88513 |
| **Reactivity** | Human |
| **Predicted** | Mouse, Rat, Bovine, Canine, Rabbit, Chicken |
| **Host** | Rabbit |
| **Clonality** | Polyclonal |
| **Isotype** | Rabbit IgG |
| **Clone Names** | RB20074 |
| **Calculated MW** | 23565 |
| **Antigen Region** | 96-123 |

## Additional Information

| | |
|---|---|
| **Gene ID** | 51053 |
| **Other Names** | Geminin, GMNN |
| **Target/Specificity** | This GMNN antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 96-123 amino acids from the Central region of human GMNN. |
| **Dilution** | WB~~1:1000 IHC-P~~1:100~500 E~~Use at an assay dependent concentration. |
| **Format** | Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification. |
| **Storage** | Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles. |
| **Precautions** | GMNN Antibody (Center Y111) is for research use only and not for use in diagnostic or therapeutic procedures. |

## Protein Information

| | |
|---|---|
| **Name** | GMNN |
| **Function** | Inhibits DNA replication by preventing the incorporation of MCM complex into pre-replication complex (pre-RC) (PubMed:14993212, PubMed:20129055, PubMed:24064211, PubMed:9635433). It is degraded during the mitotic phase |

| | |
|---|---|
| | of the cell cycle (PubMed:14993212, PubMed:24064211, PubMed:9635433). Its destruction at the metaphase- anaphase transition permits replication in the succeeding cell cycle (PubMed:14993212, PubMed:24064211, PubMed:9635433). Inhibits histone acetyltransferase activity of KAT7/HBO1 in a CDT1-dependent manner, inhibiting histone H4 acetylation and DNA replication licensing (PubMed:20129055). Inhibits the transcriptional activity of a subset of Hox proteins, enrolling them in cell proliferative control (PubMed:22615398). |
| **Cellular Location** | Cytoplasm. Nucleus. Note=Mainly cytoplasmic but can be relocalized to the nucleus. |


## Background

GMNN inhibits DNA replication by preventing the incorporation of MCM complex into prereplication complex (pre-RC). It is degraded during the mitotic phase of the cell cycle. Its destruction at the metaphase-anaphase transition permits replication in the succeeding cell cycle.


## References


De Marco, V., et al. Proc. Natl. Acad. Sci. U.S.A. 106(47):19807-19812(2009)
Zhu, W., et al. Cancer Res. 69(11):4870-4877(2009)
Lu, F., et al. Biol. Cell 101(5):273-285(2009)


## Images




Western blot analysis of GMNN Antibody (Center Y111) (Cat. #AP5227c) in HepG2 cell line lysates (35ug/lane).GMNN (arrow) was detected using the purified Pab.




GMNN Antibody (Center Y111) (Cat. #AP5227c) IHC analysis in formalin fixed and paraffin embedded lung tissue followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of the GMNN Antibody (Center Y111) for immunohistochemistry. Clinical relevance has not been evaluated.


Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.