

CD33 Rabbit pAb

CD33 Rabbit pAb
Catalog # AP52282

Product Information

Application	WB
Primary Accession	Q63994
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	44824
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from mouse CD33
Epitope Specificity	101-200/334
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SUBCELLULAR LOCATION	Cell membrane; Single-pass type I membrane protein.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	Enables protein phosphatase binding activity and sialic acid binding activity. Involved in several processes, including negative regulation of cytokine production; negative regulation of monocyte activation; and positive regulation of protein tyrosine phosphatase activity. Located in several cellular components, including Golgi apparatus; external side of plasma membrane; and peroxisome. [provided by Alliance of Genome Resources, Apr 2022]

Additional Information

Gene ID	12489
Other Names	Myeloid cell surface antigen CD33, Sialic acid-binding Ig-like lectin 3, Siglec-3, CD33, Cd33, Siglec3
Dilution	WB=1:500-2000,Flow-Cyt=1:100-200
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Protein Information

Name	Cd33
-------------	------

Synonyms	Siglec3
Function	Sialic-acid-binding immunoglobulin-like lectin (Siglec) that plays a role in mediating cell-cell interactions and in maintaining immune cells in a resting state (By similarity). Preferentially binds sialic acid to the short O-linked glycans of certain mucins (PubMed: 12773563).
Cellular Location	Cell membrane {ECO:0000250 UniProtKB:P20138}; Single-pass type I membrane protein {ECO:0000250 UniProtKB:P20138}
Tissue Location	Expressed on myeloid precursors in the bone marrow. In the peripheral blood, mostly expressed on granulocytes


Background

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

References

Tchilian E.Z.,et al.Blood 83:3188-3198(1994).

Images



Sample:

Lane 1: Bone (Mouse) Lysate at 40 ug

Lane 2: Lymph node (Mouse) Lysate at 40 ug

Primary: Anti-CD33 (AP52282) at 1/1000 dilution

Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution

Predicted band size: 40/37 kD

Observed band size: 65 kD

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.