

LYVE-1 Rabbit pAb

LYVE-1 Rabbit pAb
Catalog # AP52292

Product Information

Application	IHC-P, IHC-F, IF
Primary Accession	Q8BHC0
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	34574
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from mouse LYVE-1 terminus
Epitope Specificity	231-318/318
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SUBCELLULAR LOCATION	Membrane; Single-pass type I membrane protein. Note=Localized to the plasma membrane and in vesicles near extranuclear membranes which may represent trans-Golgi network (TGN) and endosomes/prelysosomal compartments. Undergoes ligand-dependent internalization and recycling at the cell surface.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	The lymphatic vasculature forms a second circulatory system that drains extracellular fluid from the tissues and provides an exclusive environment in which immune cells can encounter and respond to foreign antigen. Recently a number of interesting molecules have been identified that may be exploited as markers for lymphatic endothelium, including the hyaluronan receptor LYVE1, PALE, VEGFR3, podoplanin. LYVE1 has been identified as a major receptor for HA (extracellular matrix glycosaminoglycan hyaluronan) on the lymph vessel wall. The deduced amino acid sequence of LYVE1 predicts a 322-residue type I integral membrane polypeptide 41% similar to the CD44 HA receptor with a 212-residue extracellular domain containing a single Link module the prototypic HA binding domain of the Link protein superfamily. Like CD44, the LYVE1 molecule binds both soluble and immobilized HA. However, unlike CD44, the LYVE1 molecule colocalizes with HA on the luminal face of the lymph vessel wall and is completely absent from blood vessels. Hence, LYVE1 is the first lymph-specific HA receptor to be characterized and is a uniquely powerful marker for lymph vessels themselves.

Additional Information

Gene ID	114332
Other Names	Lymphatic vessel endothelial hyaluronic acid receptor 1, LYVE-1, Cell surface

retention sequence-binding protein 1, CRSBP-1, Extracellular link domain-containing protein 1, Lyve1, Crsbp1, Xlkd1

Dilution

IHC-P=1:200-500,IHC-F=1:200-500,IF=1:200-500

Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Protein Information

Name

Lyve1

Synonyms

Crsbp1, Xlkd1

Function

Ligand-specific transporter trafficking between intracellular organelles (TGN) and the plasma membrane. Plays a role in autocrine regulation of cell growth mediated by growth regulators containing cell surface retention sequence binding (CRS). May act as a hyaluronan (HA) transporter, either mediating its uptake for catabolism within lymphatic endothelial cells themselves, or its transport into the lumen of afferent lymphatic vessels for subsequent re-uptake and degradation in lymph nodes (PubMed:[10187853](#)). Binds to pericellular hyaluronan matrices deposited on the surface of leukocytes and facilitates cell adhesion and migration through lymphatic endothelium (By similarity).

Cellular Location

Membrane; Single- pass type I membrane protein Note=Localized to the plasma membrane and in vesicles near extranuclear membranes which may represent trans-Golgi network (TGN) and endosomes/prelysosomal compartments. Undergoes ligand-dependent internalization and recycling at the cell surface


Background

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

References

Banerji S.,et al.J. Cell Biol. 144:789-801(1999).
Carninci P.,et al.Science 309:1559-1563(2005).
Boensch C.,et al.J. Biol. Chem. 274:10582-10589(1999).
Bernhard O.K.,et al.J. Proteome Res. 6:987-995(2007).

Images



Paraformaldehyde-fixed, paraffin embedded Human Liver; Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15 min; The section was incubated with LYVE-1 Polyclonal Antibody, Unconjugated (AP52292) at 1:600 overnight at 4°C, followed by conjugation to the AP52292-HRP and DAB (C-0010) staining.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.