

IL-23 Rabbit pAb

IL-23 Rabbit pAb
Catalog # AP52327

Product Information

Application	WB, IHC-P, IHC-F, IF, E
Primary Accession	Q9NPF7
Reactivity	Human, Mouse
Predicted	Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	20730
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human IL-23A
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SUBCELLULAR LOCATION	Secreted. Secreted upon association with IL12B.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Background Descriptions	IL-23 is a heterodimeric cytokine composed of two subunits: a unique p19 subunit, and a p40 subunit that is shared with IL-12. IL-23 is secreted by activated dendritic cells and macrophages. IL-23 binds to a heterodimeric receptor complex composed of IL12RB1 and IL23R, activates the Jak-Stat signaling cascade, stimulates memory rather than naive T cells and promotes production of proinflammatory cytokines, such as IFNγ and IL-17. IL-23 induces autoimmune inflammation and thus may be responsible for autoimmune inflammatory diseases and may be important for tumorigenesis. This recombinant human IL-23 is produced by human cells.(Tag free) Biological activity: The activity was measured by its ability to induce the IL-17 expression in murine splenocytes. Reconstitution: Briefly centrifuge the vial before opening. It is recommended to reconstitute the protein in sterile PBS containing 0.1% endotoxin-free recombinant human serum albumin.

Additional Information

Gene ID	51561
Other Names	Interleukin-23 subunit alpha, IL-23 subunit alpha, IL-23-A, Interleukin-23 subunit p19, IL-23p19, IL23A, SGRF
Dilution	WB=1:500-2000,IHC-P=1:100-500,IHC-F=1:100-500,IF=1:100-500,ELISA=1:5000-10000
Storage	Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody

is stable for at least two weeks at 2-4 °C.

Protein Information

Name	IL23A
Synonyms	SGRF
Function	Associates with IL12B to form the pro-inflammatory cytokine IL-23 that plays different roles in innate and adaptive immunity (PubMed: 11114383). Released by antigen-presenting cells such as dendritic cells or macrophages, binds to a heterodimeric receptor complex composed of IL12RB1 and IL23R to activate JAK2 and TYK2 which then phosphorylate the receptor to form a docking site leading to the phosphorylation of STAT3 and STAT4 (PubMed: 29287995 , PubMed: 32474165 , PubMed: 33606986). This process leads to activation of several pathways including p38 MAPK or NF-kappa-B and promotes the production of pro- inflammatory cytokines such as interleukin-17A/IL17A (PubMed: 12023369). In turn, participates in the early and effective intracellular bacterial clearance (PubMed: 32474165). Promotes the expansion and survival of T-helper 17 cells, a CD4-positive helper T-cell subset that produces IL-17, as well as other IL-17-producing cells (PubMed: 17676044).
Cellular Location	Secreted. Note=Secreted upon association with IL12B
Tissue Location	Secreted by activated dendritic and phagocytic cells and keratinocytes. Also expressed by dermal Langerhans cells (at protein level).


Background

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

References

Oppmann B.,et al.Immunity 13:715-725(2000).
Hirata Y.,et al.Submitted (JUL-1999) to the EMBL/GenBank/DDBJ databases.
Clark H.F.,et al.Genome Res. 13:2265-2270(2003).
Parham C.,et al.J. Immunol. 168:5699-5708(2002).
Pirhonen J.,et al.J. Immunol. 169:5673-5678(2002).

Images



Sample:

Lane 1: Mouse Ear tissue lysates

Lane 2: Human A431 cell lysates


Lane 3: Human Jurkat cell lysates

Primary: Anti-IL-23 (AP52327) at 1/1000 dilution

Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution

Predicted band size: 21 kDa

Observed band size: 21 kDa



Tissue/cell: human rhinopolypus tissue; 4%
Paraformaldehyde-fixed and paraffin-embedded;
Antigen retrieval: citrate buffer (0.01M, pH 6.0), Boiling
bathing for 15min; Block endogenous peroxidase by 3%
Hydrogen peroxide for 30min; Blocking buffer (normal
goat serum,C-0005) at 37°C for 20 min;
Incubation: Anti-IL-23 Polyclonal Antibody,
Unconjugated(AP52327) 1:300, overnight at 4°C, followed
by conjugation to the secondary antibody(SP-0023) and
DAB(C-0010) staining

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.