

# Smad1 (Phospho-Ser465) Antibody

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP52355

## Product Information

Application	WB, IHC
Primary Accession	<a href="#">Q15797</a>
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	52260

## Additional Information

Gene ID	4086
Other Names	Mothers against decapentaplegic homolog 1, MAD homolog 1, Mothers against DPP homolog 1, JV4-1, Mad-related protein 1, SMAD family member 1, SMAD 1, Smad1, hSMAD1, Transforming growth factor-beta-signaling protein 1, BSP-1, SMAD1, BSP1, MADH1, MADR1
Dilution	WB~~1:1000 IHC~~1:50~100
Format	Rabbit IgG in phosphate buffered saline (without Mg <sup>2+</sup> and Ca <sup>2+</sup> ), pH 7.4, 150mM NaCl, 0.09% (W/V) sodium azide and 50% glycerol.
Storage Conditions	-20°C

## Protein Information

Name	SMAD1
Synonyms	BSP1, MADH1, MADR1
Function	Transcriptional modulator that plays a role in various cellular processes, including embryonic development, cell differentiation, and tissue homeostasis (PubMed: <a href="#">9335504</a> ). Upon BMP ligand binding to their receptors at the cell surface, is phosphorylated by activated type I BMP receptors (BMPRI) and associates with SMAD4 to form a heteromeric complex which translocates into the nucleus acting as transcription factor (PubMed: <a href="#">33667543</a> ). In turn, the hetero-trimeric complex recognizes cis-regulatory elements containing Smad Binding Elements (SBEs) to modulate the outcome of the signaling network (PubMed: <a href="#">33667543</a> ). SMAD1/OAZ1/PSMB4 complex mediates the degradation of the CREBBP/EP300 repressor SNIP1. Positively regulates BMP4-induced expression of odontogenic development regulator MSX1 following IPO7-mediated nuclear import (By similarity).



<b>Cellular Location</b>	Cytoplasm. Nucleus Note=Cytoplasmic in the absence of ligand. Migrates to the nucleus when complexed with SMAD4 (PubMed:15647271). Co-localizes with LEMD3 at the nucleus inner membrane (PubMed:15647271). Exported from the nucleus to the cytoplasm when dephosphorylated (By similarity) {ECO:0000250 UniProtKB:P70340, ECO:0000269 PubMed:15647271}
<b>Tissue Location</b>	Ubiquitous. Highest expression seen in the heart and skeletal muscle


## Background

Transcriptional modulator activated by BMP (bone morphogenetic proteins) type 1 receptor kinase. SMAD1 is a receptor-regulated SMAD (R-SMAD). SMAD1/OAZ1/PSMB4 complex mediates the degradation of the CREBBP/EP300 repressor SNIP1. May act synergistically with SMAD4 and YY1 in bone morphogenetic protein (BMP)-mediated cardiac-specific gene expression.


## References

Riggins G.J.,et al.Nat. Genet. 13:347-349(1996).  
Liu F.,et al.Nature 381:620-623(1996).  
Hoodless P.A.,et al.Cell 85:489-500(1996).  
Lechleider R.J.,et al.J. Biol. Chem. 271:17617-17620(1996).  
Zhang Y.,et al.Nature 383:168-172(1996).

## Images



Western blot analysis of extracts from HeLa cells treated with Serum (10%, 15min), using Smad1 (phospho-Ser465) antibody.



Immunohistochemical analysis of paraffin-embedded human heart tissue, using Smad1 (phospho-Ser465) antibody.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.