

SF1 (Phospho-Ser82) Antibody

Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP52392

Product Information

Application WB, IHC Primary Accession Q15637

Reactivity Human, Mouse, Rat

Host Rabbit
Clonality Polyclonal
Calculated MW 68330

Additional Information

Gene ID 7536

Other Names Splicing factor 1, Mammalian branch point-binding protein, BBP, mBBP,

Transcription factor ZFM1, Zinc finger gene in MEN1 locus, Zinc finger protein

162, SF1, ZFM1, ZNF162

Dilution WB~~1:1000 IHC~~1:50~100

Format Rabbit IgG in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4,

150mM NaCl, 0.09% (W/V) sodium azide and 50% glycerol.

Storage Conditions -20°C

Protein Information

Name SF1

Synonyms ZFM1, ZNF162

Function Necessary for the ATP-dependent first step of spliceosome assembly. Binds

to the intron branch point sequence (BPS) 5'-UACUAAC-3' of the pre-mRNA.

May act as transcription repressor.

Cellular Location Nucleus.

Tissue Location Detected in lung, ovary, adrenal gland, colon, kidney, muscle, pancreas,

thyroid, placenta, brain, liver and heart

Background


Necessary for the ATP-dependent first step of spliceosome assembly. Binds to the intron branch point

sequence (BPS) 5'-UACUAAC-3' of the pre-mRNA. May act as transcription repressor.


References

Arning S.,et al.RNA 2:794-810(1996). Caslini C.,et al.Genomics 42:268-277(1997). Toda T.,et al.Hum. Mol. Genet. 3:465-470(1994). Ota T.,et al.Nat. Genet. 36:40-45(2004). Taylor T.D.,et al.Nature 440:497-500(2006).

Images



Western blot analysis of extracts from HUVEC cells, treated with anisomycin (25ug/ml, 30mins), using SF1 (Phospho-Ser82) antibody.



Immunohistochemistry analysis of paraffin-embedded human brain tissue using SF1 (Phospho-Ser82) antibody

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.