

I κ B- α (Phospho-Tyr305) Antibody

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP52429

Product Information

Application	WB, IHC
Primary Accession	P25963
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal

Additional Information

Other Names	NF-kappa-B inhibitor alpha, I-kappa-B-alpha, I κ B-alpha, IkappaBalpha, Major histocompatibility complex enhancer-binding protein MAD3, NFKBIA, IKBA, MAD3, NFKBI
Dilution	WB~~1:1000 IHC~~1:50~100
Format	Rabbit IgG in phosphate buffered saline (without Mg ²⁺ and Ca ²⁺), pH 7.4, 150mM NaCl, 0.09% (W/V) sodium azide and 50% glycerol.
Storage Conditions	-20°C

Protein Information


Background

Inhibits the activity of dimeric NF-kappa-B/REL complexes by trapping REL dimers in the cytoplasm through masking of their nuclear localization signals. On cellular stimulation by immune and proinflammatory responses, becomes phosphorylated promoting ubiquitination and degradation, enabling the dimeric RELA to translocate to the nucleus and activate transcription.


References

- Haskill S.,et al.Cell 65:1281-1289(1991).
Jungnickel B.,et al.J. Exp. Med. 191:395-402(2000).
Liu B.,et al.Submitted (APR-2001) to the EMBL/GenBank/DDBJ databases.
Kalnine N.,et al.Submitted (OCT-2004) to the EMBL/GenBank/DDBJ databases.
Ota T.,et al.Nat. Genet. 36:40-45(2004).

Images



Western blot analysis of extracts from COS7 cells, treated with nocodazole (1ug/ml, 16hours), using IkB- α (Phospho-Tyr305) antibody.



Immunohistochemistry analysis of paraffin-embedded human lymph node tissue using IkB- α (Phospho-Tyr305) antibody.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.