

IGF2R (Phospho-Ser2409) Antibody

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP52445

Product Information

Application	WB, IHC
Primary Accession	P11717
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	274375

Additional Information

Gene ID	3482
Other Names	Cation-independent mannose-6-phosphate receptor, CI Man-6-P receptor, CI-MPR, M6PR, 300 kDa mannose 6-phosphate receptor, MPR 300, Insulin-like growth factor 2 receptor, Insulin-like growth factor II receptor, IGF-II receptor, M6P/IGF2 receptor, M6P/IGF2R, CD222, IGF2R, MPRI
Dilution	WB~~1:1000 IHC~~1:50~100
Format	Rabbit IgG in phosphate buffered saline (without Mg ²⁺ and Ca ²⁺), pH 7.4, 150mM NaCl, 0.09% (W/V) sodium azide and 50% glycerol.
Storage Conditions	-20°C

Protein Information

Name	IGF2R
Synonyms	MPRI
Function	Mediates the transport of phosphorylated lysosomal enzymes from the Golgi complex and the cell surface to lysosomes (PubMed: 18817523 , PubMed: 2963003). Lysosomal enzymes bearing phosphomannosyl residues bind specifically to mannose-6-phosphate receptors in the Golgi apparatus and the resulting receptor-ligand complex is transported to an acidic prelysosomal compartment where the low pH mediates the dissociation of the complex (PubMed: 18817523 , PubMed: 2963003). The receptor is then recycled back to the Golgi for another round of trafficking through its binding to the retromer (PubMed: 18817523). This receptor also binds IGF2 (PubMed: 18046459). Acts as a positive regulator of T-cell coactivation by binding DPP4 (PubMed: 10900005).
Cellular Location	Golgi apparatus membrane; Single-pass type I membrane protein. Endosome

membrane; Single-pass type I membrane protein. Note=Mainly localized in the Golgi at steady state and not detectable in lysosome (PubMed:18817523) Colocalized with DPP4 in internalized cytoplasmic vesicles adjacent to the cell surface (PubMed:10900005).


Background

Transport of phosphorylated lysosomal enzymes from the Golgi complex and the cell surface to lysosomes. Lysosomal enzymes bearing phosphomannosyl residues bind specifically to mannose-6- phosphate receptors in the Golgi apparatus and the resulting receptor-ligand complex is transported to an acidic prelysosomal compartment where the low pH mediates the dissociation of the complex. This receptor also binds IGF2. Acts as a positive regulator of T-cell coactivation, by binding DPP4.


References

Morgan D.O.,et al.Nature 329:301-307(1987).
Oshima A.,et al.J. Biol. Chem. 263:2553-2562(1988).
Gemma A.,et al.Submitted (NOV-1998) to the EMBL/GenBank/DDBJ databases.
Killian J.K.,et al.Mamm. Genome 10:74-77(1999).
Mungall A.J.,et al.Nature 425:805-811(2003).

Images



Western blot analysis of extracts from COS-7 cells, treated with UV (15mins), using IGF2R (Phospho-Ser2409) antibody.



Immunohistochemistry analysis of paraffin-embedded human brain tissue using IGF2R (Phospho-Ser2409) antibody.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.