

POU5F1/OCT4 Antibody

Purified Mouse Monoclonal Antibody (Mab) Catalog # AP52657

Product Information

Application WB
Primary Accession Q01860
Reactivity Mouse
Host Mouse
Clonality Monoclonal
Isotype IgG2b
Calculated MW 38571

Additional Information

Gene ID 5460

Other Names MGC22487;Oct 3;Oct 4;Oct-3;Oct-4;OCT3;Oct4;Octamer binding protein

3;Octamer binding protein 4;Octamer binding transcription factor

3;Octamer-binding protein 3;Octamer-binding protein 4;Octamer-binding transcription factor 3;OTF 3;OTF 4;OTF-3;OTF3;OTF4; PO5F1_HUMAN;POU class 5 homeobox 1;POU domain class 5 transcription factor 1;POU domain transcription factor OCT4;POU domain, class 5, transcription factor 1;POU5F1.

Dilution WB~~1:1000

Format Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide, pH

7.3.

Storage Store at 4°C short term. Aliquot and store at -20°C long term. Avoid

freeze/thaw cycles.

Protein Information

Name POU5F1

Synonyms OCT3, OCT4, OTF3

Function Transcription factor that binds to the octamer motif (5'- ATTTGCAT-3'). Forms

a trimeric complex with SOX2 or SOX15 on DNA and controls the expression of a number of genes involved in embryonic development such as YES1, FGF4, UTF1 and ZFP206. Critical for early embryogenesis and for embryonic stem

cell pluripotency.

Cellular Location Cytoplasm. Nucleus. Note=Expressed in a diffuse and slightly punctuate

pattern. Colocalizes with MAPK8 and MAPK9 in the nucleus.

{ECO:0000250|UniProtKB:P20263, ECO:0000269|PubMed:18191611,

ECO:0000269 | PubMed:19274063, ECO:0000269 | PubMed:23024368 }

Tissue Location

Expressed in developing brain. Highest levels found in specific cell layers of the cortex, the olfactory bulb, the hippocampus and the cerebellum. Low levels of expression in adult tissues.


Background

Transcription factor that binds to the octamer motif (5'-ATTTGCAT-3'). Forms a trimeric complex with SOX2 on DNA and controls the expression of a number of genes involved in embryonic development such as YES1, FGF4, UTF1 and ZFP206. Critical for early embryogenesis and for embryonic stem cell pluripotency.

References

Takeda J., et al. Nucleic Acids Res. 20:4613-4620(1992). Stuart P.E., et al. Tissue Antigens 76:387-397(2010). Shiina T., et al. Genetics 173:1555-1570(2006). Shiina S., et al. Submitted (SEP-1999) to the EMBL/GenBank/DDBJ databases. Mungall A.J., et al. Nature 425:805-811(2003).

Images



Western blot detection of POU5F1/OCT4 in F9 cell lysates using POU5F1/OCT4 mouse mAb (1:1000 diluted). Predicted band size: 45KDa. Observed band size: 45KDa.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.