

Hsp70 (C-terminus) Antibody

Purified Mouse Monoclonal Antibody (Mab) Catalog # AP52677

Product Information

Application	WB, IP
Primary Accession	<u>P08107</u>
Reactivity	Human
Host	Mouse
Clonality	Monoclonal
Isotype	IgG2b
Calculated MW	70 KDa

Additional Information

Other Names	DAQB 147D11.1 001;FLJ54303;FLJ54370;FLJ54392;FLJ54408;FLJ75127;Heat shock 70 kDa protein 1;Heat shock 70 kDa protein 1/2;Heat shock 70 kDa protein 1A/1B;heat shock 70kDa protein 1A;Heat shock 70kDa protein 1B;Heat shock induced protein;heat shock protein 70;HSP70 1;HSP70 2;HSP70-1/HSP7 0-2;HSP70-1A;HSP70.1;HSP70.1/HSP70.2;HSP70I;HSP71_HUMAN;HSP72; HSPA1;HSPA1A;HSPA1B;XXbac BCX40G17.3 001.
Dilution	WB~~1:1000 IP~~1:500
Format	Purified mouse monoclonal in buffer containing 0.1M Tris-Glycine(pH 7.4,150 mM NaCl)with 0.09% (W/V) sodium azide,0.1mg/mlBSA and 50% glycerol.
Storage	Store at -20 °C.Stable for 12 months from date of receipt

Protein Information

Background

In cooperation with other chaperones, Hsp70s stabilize preexistent proteins against aggregation and mediate the folding of newly translated polypeptides in the cytosol as well as within organelles. These chaperones participate in all these processes through their ability to recognize nonnative conformations of other proteins. They bind extended peptide segments with a net hydrophobic character exposed by polypeptides during translation and membrane translocation, or following stress-induced damage. In case of rotavirus A infection, serves as a post-attachment receptor for the virus to facilitate entry into the cell.

References

Milner C.M., et al.Immunogenetics 32:242-251(1990). Hunt C., et al.Proc. Natl. Acad. Sci. U.S.A. 82:6455-6459(1985). Xie T.,et al.Genome Res. 13:2621-2636(2003). Shiina S.,et al.Submitted (SEP-1999) to the EMBL/GenBank/DDBJ databases. Ota T.,et al.Nat. Genet. 36:40-45(2004).

Images



Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.