

# **GAPDH** Antibody

Purified Mouse Monoclonal Antibody (Mab) Catalog # AP52680

### **Product Information**

Application	WB
Primary Accession	<u>P04406</u>
Reactivity	Rat, Human, Mouse, Hamster, Monkey
Host	Mouse
Clonality	Monoclonal
Isotype	IgM
Calculated MW	36053

## **Additional Information**

Gene ID	2597
Other Names	38 kDa BFA-dependent ADP-ribosylation substrate; aging associated gene 9 protein; Aging-associated gene 9 protein; BARS-38; cb609; EC 1.2.1.12; G3P_HUMAN; G3PD; G3PDH; GAPD; GAPDH; GAPDH; Glyceraldehyde 3 phosphate dehydrogenase; Glyceraldehyde 3 phosphate dehydrogenase liver; Glyceraldehyde 3 phosphate dehydrogenase muscle; Glyceraldehyde-3-phosphate dehydrogenase; KNC-NDS6; MGC102544; MGC102546; MGC103190; MGC103191; MGC105239; MGC127711; MGC88685; OCAS, p38 component; OCT1 coactivator in S phase, 38-KD component; peptidyl cysteine S nitrosylase GAPDH; Peptidyl-cysteine S-nitrosylase GAPDH; wu:fb33a10.
Dilution	WB~~1:5000
Format	Ascites containing with 0.09% (W/V) sodium azide and 50% glycerol.
Storage	Store at -20 °C.Stable for 12 months from date of receipt

#### **Protein Information**

Name	GAPDH {ECO:0000303 PubMed:2987855, ECO:0000312 HGNC:HGNC:4141}
Function	Has both glyceraldehyde-3-phosphate dehydrogenase and nitrosylase activities, thereby playing a role in glycolysis and nuclear functions, respectively (PubMed: <u>11724794</u> , PubMed: <u>3170585</u> ). Glyceraldehyde-3-phosphate dehydrogenase is a key enzyme in glycolysis that catalyzes the first step of the pathway by converting D- glyceraldehyde 3-phosphate (G3P) into 3-phospho-D-glyceroyl phosphate (PubMed: <u>11724794</u> , PubMed: <u>3170585</u> ). Modulates the organization and assembly of the cytoskeleton (By similarity). Facilitates the CHP1- dependent microtubule and membrane associations through its ability to stimulate the binding of CHP1 to

	microtubules (By similarity). Component of the GAIT (gamma interferon-activated inhibitor of translation) complex which mediates interferon-gamma-induced transcript-selective translation inhibition in inflammation processes (PubMed:23071094). Upon interferon-gamma treatment assembles into the GAIT complex which binds to stem loop-containing GAIT elements in the 3'-UTR of diverse inflammatory mRNAs (such as ceruplasmin) and suppresses their translation (PubMed:23071094). Also plays a role in innate immunity by promoting TNF-induced NF-kappa-B activation and type I interferon production, via interaction with TRAF2 and TRAF3, respectively (PubMed:2332158, PubMed:27387501). Participates in nuclear events including transcription, RNA transport, DNA replication and apoptosis (By similarity). Nuclear functions are probably due to the nitrosylase activity that mediates cysteine S-nitrosylation of nuclear target proteins such as SIRT1, HDAC2 and PRKDC (By similarity).
Cellular Location	Cytoplasm, cytosol. Nucleus {ECO:0000250 UniProtKB:P04797}. Cytoplasm, perinuclear region. Membrane Cytoplasm, cytoskeleton {ECO:0000250 UniProtKB:P04797} Note=Translocates to the nucleus following S-nitrosylation and interaction with SIAH1, which contains a nuclear localization signal (By similarity). Postnuclear and Perinuclear regions (PubMed:12829261) {ECO:0000250 UniProtKB:P04797, ECO:0000269 PubMed:12829261}

### Background

Has both glyceraldehyde-3-phosphate dehydrogenase and nitrosylase activities, thereby playing a role in glycolysis and nuclear functions, respectively. Participates in nuclear events including transcription, RNA transport, DNA replication and apoptosis. Nuclear functions are probably due to the nitrosylase activity that mediates cysteine S-nitrosylation of nuclear target proteins such as SIRT1, HDAC2 and PRKDC. Modulates the organization and assembly of the cytoskeleton. Facilitates the CHP1-dependent microtubule and membrane associations through its ability to stimulate the binding of CHP1 to microtubules (By similarity). Glyceraldehyde-3-phosphate dehydrogenase is a key enzyme in glycolysis that catalyzes the first step of the pathway by converting D-glyceraldehyde 3-phosphate (G3P) into 3-phospho-D- glyceroyl phosphate. Component of the GAIT (gamma interferon- activated inhibitor of translation) complex which mediates interferon-gamma-induced transcript-selective translation inhibition in inflammation processes. Upon interferon-gamma treatment assembles into the GAIT complex which binds to stem loop-containing GAIT elements in the 3'-UTR of diverse inflammatory mRNAs (such as ceruplasmin) and suppresses their translation.

#### References

Hanauer A., et al.EMBO J. 3:2627-2633(1984). Arcari P., et al.Nucleic Acids Res. 12:9179-9189(1984). Tso J.Y., et al.Nucleic Acids Res. 13:2485-2502(1985). Tokunaga K., et al.Cancer Res. 47:5616-5619(1987). Allen R.W., et al.J. Biol. Chem. 262:649-653(1987).

#### Images



Western blot detection of GAPDH in Hela,3T3,C6,COS7,CHO-K1,Rat brain and Jurkat cell lysates using GAPDH mouse mAb (1:5000 diluted).Predicted band size:36KDa.Observed band size:36KDa. Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.