

Cyclin H Antibody

Purified Mouse Monoclonal Antibody (Mab)

Catalog # AP52723

Product Information

Application	WB, IP
Primary Accession	P51946
Reactivity	Human
Host	Mouse
Clonality	Monoclonal
Isotype	IgG2b
Calculated MW	37643

Additional Information

Gene ID	902
Other Names	6330408H09Rik;AI661354;AV102684;AW538719;CAK;CAK complex subunit;ccnh;CCNH_HUMAN;CDK activating kinase;CDK activating kinase complex subunit;Cyclin dependent kinase activating kinase;cyclin dependent kinase activating kinase complex subunit;Cyclin H; CyclinH;MO15-associated protein;p34;p36;p37.
Dilution	WB~~1:1000 IP~~1:500
Format	Purified mouse monoclonal in buffer containing 0.1M Tris-Glycine (pH 7.4, 150 mM NaCl) with 0.09% (W/V) sodium azide, 50% glycerol
Storage	Store at -20 °C.Stable for 12 months from date of receipt

Protein Information

Name	CCNH
Function	Regulates CDK7, the catalytic subunit of the CDK-activating kinase (CAK) enzymatic complex. CAK activates the cyclin-associated kinases CDK1, CDK2, CDK4 and CDK6 by threonine phosphorylation. CAK complexed to the core-TFIID basal transcription factor activates RNA polymerase II by serine phosphorylation of the repetitive C-terminal domain (CTD) of its large subunit (POLR2A), allowing its escape from the promoter and elongation of the transcripts. Involved in cell cycle control and in RNA transcription by RNA polymerase II. Its expression and activity are constant throughout the cell cycle.
Cellular Location	Nucleus.


Background

Regulates CDK7, the catalytic subunit of the CDK- activating kinase (CAK) enzymatic complex. CAK activates the cyclin-associated kinases CDK1, CDK2, CDK4 and CDK6 by threonine phosphorylation. CAK complexed to the core-TFIIF basal transcription factor activates RNA polymerase II by serine phosphorylation of the repetitive C-terminal domain (CTD) of its large subunit (POLR2A), allowing its escape from the promoter and elongation of the transcripts. Involved in cell cycle control and in RNA transcription by RNA polymerase II. Its expression and activity are constant throughout the cell cycle.


References

Maekela T.P.,et al.Nature 371:254-257(1994).
Fisher R.P.,et al.Cell 78:713-724(1994).
Ebert L.,et al.Submitted (MAY-2004) to the EMBL/GenBank/DDBJ databases.
Shiekhhattar R.,et al.Nature 374:283-287(1995).
Kershner E.,et al.J. Biol. Chem. 273:34444-34453(1998).

Images



Western blot detection of Cyclin H in Jurkat,K562 and A431 cell lysates using Cyclin H mouse mAb (1:1000 diluted).Predicted band size: 38KDa.Observed band size: 38KDa.



Immunoprecipitation analysis of HeLa cell lysates using Cyclin H mouse mAb.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.