

GRK2 Antibody

Purified Mouse Monoclonal Antibody (Mab)

Catalog # AP52726

Product Information

Primary Accession	P25098
Host	Mouse
Clonality	Monoclonal
Isotype	IgG1
Calculated MW	79574

Additional Information

Gene ID	156
Other Names	ADRBK 1;ADRBK1;Adrenergic beta receptor kinase 1;Adrenergic beta receptor kinase 1; ARBK1_HUMAN;BARK 1;BARK;BARK1;Beta adrenergic receptor kinase 1;Beta ARK 1;Beta ARK1; Beta-adrenergic receptor kinase 1;Beta-ARK-1;FLJ16718;G Protein Coupled Receptor Kinase 2;G protein dependent receptor kinase 2;G-protein coupled receptor kinase 2;GRK 2;GRK2.
Format	Purified mouse monoclonal in buffer containing 0.1M Tris-Glycine (pH 7.4, 150 mM NaCl) with 0.09% (W/V) sodium azide, 50% glycerol
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.

Protein Information

Name	GRK2 {ECO:0000303 PubMed:40384633, ECO:0000312 HGNC:HGNC:289}
Function	Specifically phosphorylates the activated forms of G protein- coupled receptors (PubMed: 14976224 , PubMed: 16179383 , PubMed: 19306925 , PubMed: 19715378 , PubMed: 21799898 , PubMed: 23077507 , PubMed: 30284927 , PubMed: 39138140 , PubMed: 40384633). Such receptor phosphorylation initiates beta-arrestin-mediated receptor internalization, and signaling events leading to their desensitization (PubMed: 14976224 , PubMed: 16179383 , PubMed: 19306925 , PubMed: 19715378 , PubMed: 21799898 , PubMed: 23077507 , PubMed: 30284927 , PubMed: 39138140 , PubMed: 40384633). Phosphorylates the agonist-occupied form of the beta-adrenergic and closely related receptors, probably inducing a desensitization of them (PubMed: 19715378). Phosphorylates catecholamine-activated ADRB2 to regulate physiological cardiomyocyte contraction rate responses (By similarity). Also phosphorylates ligand-bound C3a and C5a anaphylatoxin receptors (C3AR1 and C5AR1, respectively), leading to receptor desensitization (PubMed: 21799898 , PubMed: 23077507).

Positively regulates ciliary smoothened (SMO) signaling pathway by mediating phosphorylation of SMO, facilitating SMO trafficking into the cilium and the stimulation of SMO activity (PubMed:[39138140](#)). Also positively regulates SMO signaling by mediating phosphorylation of GPR161, a key inhibitor SMO signaling, promoting ciliary export of GPR161 (PubMed:[40384633](#)). Key regulator of LPAR1 signaling (PubMed:[19306925](#)). Competes with RALA for binding to LPAR1 thus affecting the signaling properties of the receptor (PubMed:[19306925](#)). Desensitizes LPAR1 and LPAR2 in a phosphorylation-independent manner (PubMed:[19306925](#)). Inhibits relaxation of airway smooth muscle in response to blue light (PubMed:[30284927](#)).

Cellular Location

Cytoplasm {ECO:0000250|UniProtKB:P26817}. Cell membrane {ECO:0000250|UniProtKB:P21146}. Cell projection, cilium membrane. Postsynapse {ECO:0000250|UniProtKB:P26817}. Presynapse {ECO:0000250|UniProtKB:P26817}

Tissue Location

Expressed in peripheral blood leukocytes.


Background

Specifically phosphorylates the agonist-occupied form of the beta-adrenergic and closely related receptors, probably inducing a desensitization of them. Key regulator of LPAR1 signaling. Competes with RALA for binding to LPAR1 thus affecting the signaling properties of the receptor. Desensitizes LPAR1 and LPAR2 in a phosphorylation-independent manner.

References

Benovic J.L.,et al.FEBS Lett. 283:122-126(1991).
 Chuang T.T.,et al.J. Biol. Chem. 267:6886-6892(1992).
 Penn R.B.,et al.J. Biol. Chem. 269:14924-14930(1994).
 Aragay A.M.,et al.Proc. Natl. Acad. Sci. U.S.A. 95:2985-2990(1998).
 Oppermann M.,et al.J. Biol. Chem. 274:8875-8885(1999).

Images



Western blot detection of GRK2 in Ramos and THP-1 cell lysates using GRK2 mouse mAb (1:500 diluted),with super ECL.Predicted band size:80KDa.Observed band size:80KDa.Exposure time:20S.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.