

# eEF2 Antibody

Purified Mouse Monoclonal Antibody (Mab)

Catalog # AP52728

## Product Information

---

Application	WB, ICC
Primary Accession	<a href="#">P13639</a>
Reactivity	Human
Host	Mouse
Clonality	Monoclonal
Isotype	IgG2b
Calculated MW	95338

## Additional Information

---

Gene ID	1938
Other Names	EEF2;Eef2;EF-2;EF2;EF2_HUMAN;Elongation factor 2;Eukaryotic translation elongation factor 2;Polypeptidyl tRNA translocase;SCA26.
Dilution	WB~~1:5000 ICC~~1:200
Format	Purified mouse monoclonal in buffer containing 0.1M Tris-Glycine (pH 7.4, 150 mM NaCl) with 0.09% (W/V) sodium azide, 50%,glycerol
Storage	Store at -20 °C.Stable for 12 months from date of receipt

## Protein Information

---

Name	EEF2
Synonyms	EF2
Function	Catalyzes the GTP-dependent ribosomal translocation step during translation elongation (PubMed: <a href="#">26593721</a> ). During this step, the ribosome changes from the pre-translocational (PRE) to the post- translocational (POST) state as the newly formed A-site-bound peptidyl- tRNA and P-site-bound deacylated tRNA move to the P and E sites, respectively (PubMed: <a href="#">26593721</a> ). Catalyzes the coordinated movement of the two tRNA molecules, the mRNA and conformational changes in the ribosome (PubMed: <a href="#">26593721</a> ).
Cellular Location	Cytoplasm. Nucleus. Note=Phosphorylation by CSK promotes cleavage and SUMOylation-dependent nuclear translocation of the C- terminal cleavage product.


## Background

Catalyzes the GTP-dependent ribosomal translocation step during translation elongation. During this step, the ribosome changes from the pre-translocational (PRE) to the post- translocational (POST) state as the newly formed A-site-bound peptidyl-tRNA and P-site-bound deacylated tRNA move to the P and E sites, respectively. Catalyzes the coordinated movement of the two tRNA molecules, the mRNA and conformational changes in the ribosome.


## References

Rapp G.,et al.Biol. Chem. Hoppe-Seyler 370:1071-1075(1989).  
Hanes J.,et al.Biol. Chem. Hoppe-Seyler 373:201-204(1992).  
Ustek D.,et al.Submitted (FEB-2005) to the EMBL/GenBank/DDBJ databases.  
Mural R.J.,et al.Submitted (SEP-2005) to the EMBL/GenBank/DDBJ databases.  
Rapp G.,et al.Biol. Chem. Hoppe-Seyler 369:247-250(1988).

## Images



Western blot detection of eEF2 in HL-60,Jurkat,SHSY-5Y,U2OS and Hela cell lysates using eEF2 mouse mAb (1:5000 diluted).Predicted band size:95KDa.Observed band size: 95KDa.Exposure time:15s.



Immunocytochemistry staining of HeLa cells fixed with -20°C Methanol and using anti-eEF2 mouse mAb (dilution 1:200).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.