

c-JUN Antibody

Purified Mouse Monoclonal Antibody (Mab)

Catalog # AP52741

Product Information

Application	WB
Primary Accession	P05412
Reactivity	Transfected
Host	Mouse
Clonality	Monoclonal
Isotype	IgG1
Calculated MW	35676

Additional Information

Gene ID	3725
Other Names	Activator Protein 1;AP 1;AP1;cJun;Enhancer Binding Protein AP1;Jun Activation Domain Binding Protein;JUN;Jun oncogene;JUN protein;Jun proto oncogene;JUN_HUMAN;JUNC;Oncogene JUN;p39;Proto oncogene c jun;Proto oncogene cJun;Proto-oncogene c-jun;Transcription Factor AP 1;Transcription factor AP-1;Transcription Factor AP1;V jun avian sarcoma virus 17 oncogene homolog;V jun sarcoma virus 17 oncogene homolog (avian);V jun sarcoma virus 17 oncogene homolog;V-jun avian sarcoma virus 17 oncogene homolog;vJun Avian Sarcoma Virus 17 Oncogene Homolog.
Dilution	WB~~1:1000
Format	Purified mouse monoclonal in PBS(pH 7.4) containing with 0.09% (W/V) sodium azide and 50% glycerol.
Storage	Store at -20 °C.Stable for 12 months from date of receipt

Protein Information

Name	JUN
Function	Transcription factor that recognizes and binds to the AP-1 consensus motif 5'-TGA[GC]TCA-3' (PubMed: 10995748 , PubMed: 22083952). Heterodimerizes with proteins of the FOS family to form an AP-1 transcription complex, thereby enhancing its DNA binding activity to the AP-1 consensus sequence 5'-TGA[GC]TCA-3' and enhancing its transcriptional activity (By similarity). Together with FOSB, plays a role in activation-induced cell death of T cells by binding to the AP-1 promoter site of FASLG/CD95L, and inducing its transcription in response to activation of the TCR/CD3 signaling pathway (PubMed: 12618758). Promotes activity of NR5A1 when phosphorylated by HIPK3 leading to increased steroidogenic gene expression upon cAMP

signaling pathway stimulation (PubMed:[17210646](#)). Involved in activated KRAS-mediated transcriptional activation of USP28 in colorectal cancer (CRC) cells (PubMed:[24623306](#)). Binds to the USP28 promoter in colorectal cancer (CRC) cells (PubMed:[24623306](#)).

Cellular Location

Nucleus.

Tissue Location

Expressed in the developing and adult prostate and prostate cancer cells.


Background

Transcription factor that recognizes and binds to the enhancer heptamer motif 5'-TGA[CG]TCA-3'. Promotes activity of NR5A1 when phosphorylated by HIPK3 leading to increased steroidogenic gene expression upon cAMP signaling pathway stimulation.

References

Hattori K.,et al.Proc. Natl. Acad. Sci. U.S.A. 85:9148-9152(1988).
Bohmann D.,et al.Science 238:1386-1392(1987).
Ebert L.,et al.Submitted (JUN-2004) to the EMBL/GenBank/DDBJ databases.
Kalnine N.,et al.Submitted (OCT-2004) to the EMBL/GenBank/DDBJ databases.
Gregory S.G.,et al.Nature 441:315-321(2006).

Images



Western blot detection of c-JUN in CHO-K1 transfected by PEGFP-C1-c-JUN(A)and CHO-K1 cell lysate(B)cell lysate using c-JUN mouse mAb (1:1000 diluted).Predicted band size:43/48KDa.Observed band size:43/48KDa.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.