

# Ku70 Antibody

Purified Mouse Monoclonal Antibody (Mab)  
Catalog # AP52823

## Product Information

---

<b>Application</b>	WB, ICC
<b>Primary Accession</b>	<a href="#">P12956</a>
<b>Reactivity</b>	Human
<b>Host</b>	Mouse
<b>Clonality</b>	Monoclonal
<b>Isotype</b>	IgG2b
<b>Calculated MW</b>	69843

## Additional Information

---

<b>Gene ID</b>	2547
<b>Other Names</b>	5"-deoxyribose-5-phosphate lyase Ku70;5"-dRP lyase Ku70;70 kDa subunit of Ku antigen;ATP dependent DNA helicase 2 subunit 1;ATP dependent DNA helicase II 70 kDa subunit;ATP-dependent DNA helicase 2 subunit 1;ATP-dependent DNA helicase II 70 kDa subunit;CTC box binding factor 75 kDa subunit;CTC box-binding factor 75 kDa subunit;CTC75;CTCBF;CTCBF;DNA repair protein XRCC6;G22P1;Ku 70;Ku autoantigen 70kDa;Ku autoantigen p70 subunit;Ku autoantigen, 70kDa;Ku p70;Ku70;Ku70 DNA binding component of DNA-dependent proteinkinase complex (thyroid autoantigen 70 kDa;Kup70;Lupus Ku autoantigen protein p70;ML8;Thyroid autoantigen 70kD (Ku antigen);Thyroid autoantigen;Thyroid lupus autoantigen;Thyroid lupus autoantigen;Thyroid lupus autoantigen p70;Thyroid-lupus autoantigen;TLAA;TLAA;X ray repair complementing defective repair in Chinese hamster cells 6;X-ray repair complementing defective repair in Chinese hamster cells 6;X-ray repair cross-complementing protein 6;XRCC 6;XRCC6;XRCC6_HUMAN.
<b>Dilution</b>	WB~~1:1000 ICC~~1:200
<b>Format</b>	Purified mouse monoclonal antibody in PBS(pH 7.4) containing with 0.09% (W/V) sodium azide and 50% glycerol.
<b>Storage</b>	Store at -20 °C.Stable for 12 months from date of receipt

## Protein Information

---

<b>Name</b>	XRCC6
<b>Synonyms</b>	G22P1
	Single-stranded DNA-dependent ATP-dependent helicase that plays a key

<b>Function</b>	role in DNA non-homologous end joining (NHEJ) by recruiting DNA-PK to DNA (PubMed: <a href="#">11493912</a> , PubMed: <a href="#">12145306</a> , PubMed: <a href="#">20493174</a> , PubMed: <a href="#">2466842</a> , PubMed: <a href="#">7957065</a> , PubMed: <a href="#">8621488</a> , PubMed: <a href="#">9742108</a> ). Required for double-strand break repair and V(D)J recombination (PubMed: <a href="#">11493912</a> , PubMed: <a href="#">12145306</a> , PubMed: <a href="#">20493174</a> , PubMed: <a href="#">2466842</a> , PubMed: <a href="#">7957065</a> , PubMed: <a href="#">8621488</a> , PubMed: <a href="#">9742108</a> ). Also has a role in chromosome translocation (PubMed: <a href="#">11493912</a> , PubMed: <a href="#">12145306</a> , PubMed: <a href="#">20493174</a> , PubMed: <a href="#">2466842</a> , PubMed: <a href="#">7957065</a> , PubMed: <a href="#">8621488</a> , PubMed: <a href="#">9742108</a> ). Has a role in chromosome translocation (PubMed: <a href="#">11493912</a> , PubMed: <a href="#">12145306</a> , PubMed: <a href="#">20493174</a> , PubMed: <a href="#">2466842</a> , PubMed: <a href="#">7957065</a> , PubMed: <a href="#">8621488</a> , PubMed: <a href="#">9742108</a> ). The DNA helicase II complex binds preferentially to fork-like ends of double-stranded DNA in a cell cycle-dependent manner (PubMed: <a href="#">11493912</a> , PubMed: <a href="#">12145306</a> , PubMed: <a href="#">20493174</a> , PubMed: <a href="#">2466842</a> , PubMed: <a href="#">7957065</a> , PubMed: <a href="#">8621488</a> , PubMed: <a href="#">9742108</a> ). It works in the 3'-5' direction (PubMed: <a href="#">11493912</a> , PubMed: <a href="#">12145306</a> , PubMed: <a href="#">20493174</a> , PubMed: <a href="#">2466842</a> , PubMed: <a href="#">7957065</a> , PubMed: <a href="#">8621488</a> , PubMed: <a href="#">9742108</a> ). During NHEJ, the XRCC5-XRRC6 dimer performs the recognition step: it recognizes and binds to the broken ends of the DNA and protects them from further resection (PubMed: <a href="#">11493912</a> , PubMed: <a href="#">12145306</a> , PubMed: <a href="#">20493174</a> , PubMed: <a href="#">2466842</a> , PubMed: <a href="#">7957065</a> , PubMed: <a href="#">8621488</a> , PubMed: <a href="#">9742108</a> ). Binding to DNA may be mediated by XRCC6 (PubMed: <a href="#">11493912</a> , PubMed: <a href="#">12145306</a> , PubMed: <a href="#">20493174</a> , PubMed: <a href="#">2466842</a> , PubMed: <a href="#">7957065</a> , PubMed: <a href="#">8621488</a> , PubMed: <a href="#">9742108</a> ). The XRCC5-XRRC6 dimer acts as a regulatory subunit of the DNA-dependent protein kinase complex DNA-PK by increasing the affinity of the catalytic subunit PRKDC to DNA by 100-fold (PubMed: <a href="#">11493912</a> , PubMed: <a href="#">12145306</a> , PubMed: <a href="#">20493174</a> , PubMed: <a href="#">2466842</a> , PubMed: <a href="#">7957065</a> , PubMed: <a href="#">8621488</a> , PubMed: <a href="#">9742108</a> ). The XRCC5-XRRC6 dimer is probably involved in stabilizing broken DNA ends and bringing them together (PubMed: <a href="#">11493912</a> , PubMed: <a href="#">12145306</a> , PubMed: <a href="#">20493174</a> , PubMed: <a href="#">2466842</a> , PubMed: <a href="#">7957065</a> , PubMed: <a href="#">8621488</a> , PubMed: <a href="#">9742108</a> ). The assembly of the DNA-PK complex to DNA ends is required for the NHEJ ligation step (PubMed: <a href="#">11493912</a> , PubMed: <a href="#">12145306</a> , PubMed: <a href="#">20493174</a> , PubMed: <a href="#">2466842</a> , PubMed: <a href="#">7957065</a> , PubMed: <a href="#">8621488</a> , PubMed: <a href="#">9742108</a> ). Probably also acts as a 5'-deoxyribose-5-phosphate lyase (5'-dRP lyase), by catalyzing the beta-elimination of the 5' deoxyribose-5-phosphate at an abasic site near double-strand breaks (PubMed: <a href="#">20383123</a> ). 5'-dRP lyase activity allows to 'clean' the termini of abasic sites, a class of nucleotide damage commonly associated with strand breaks, before such broken ends can be joined (PubMed: <a href="#">20383123</a> ). The XRCC5-XRRC6 dimer together with APEX1 acts as a negative regulator of transcription (PubMed: <a href="#">8621488</a> ). In association with NAA15, the XRCC5-XRRC6 dimer binds to the osteocalcin promoter and activates osteocalcin expression (PubMed: <a href="#">12145306</a> ). Plays a role in the regulation of DNA virus-mediated innate immune response by assembling into the HDP-RNP complex, a complex that serves as a platform for IRF3 phosphorylation and subsequent innate immune response activation through the cGAS-STING pathway (PubMed: <a href="#">28712728</a> ). Negatively regulates apoptosis by interacting with BAX and sequestering it from the mitochondria (PubMed: <a href="#">15023334</a> ). Might have deubiquitination activity, acting on BAX (PubMed: <a href="#">18362350</a> ).
<b>Cellular Location</b>	Nucleus. Chromosome. Cytoplasm. Note=When trimethylated, localizes in the cytoplasm.

## Background


Single-stranded DNA-dependent ATP-dependent helicase. Has a role in chromosome translocation. The DNA helicase II complex binds preferentially to fork-like ends of double-stranded DNA in a cell cycle-dependent manner. It works in the 3'-5' direction. Binding to DNA may be mediated by XRCC6.

Involved in DNA non-homologous end joining (NHEJ) required for double-strand break repair and V(D)J recombination. The XRCC5/6 dimer acts as regulatory subunit of the DNA-dependent protein kinase complex DNA-PK by increasing the affinity of the catalytic subunit PRKDC to DNA by 100-fold. The XRCC5/6 dimer is probably involved in stabilizing broken DNA ends and bringing them together. The assembly of the DNA-PK complex to DNA ends is required for the NHEJ ligation step. Required for osteocalcin gene expression. Probably also acts as a 5'-deoxyribose-5-phosphate lyase (5'-dRP lyase), by catalyzing the beta-elimination of the 5' deoxyribose- 5-phosphate at an abasic site near double-strand breaks. 5'-dRP lyase activity allows to 'clean' the termini of abasic sites, a class of nucleotide damage commonly associated with strand breaks, before such broken ends can be joined. The XRCC5/6 dimer together with APEX1 acts as a negative regulator of transcription.


## References

- Chan J.Y.,et al.J. Biol. Chem. 264:3651-3654(1989).  
Reeves W.H.,et al.J. Biol. Chem. 264:5047-5052(1989).  
Griffith A.J.,et al.Mol. Biol. Rep. 16:91-97(1992).  
Ota T.,et al.Nat. Genet. 36:40-45(2004).  
Halleck A.,et al.Submitted (JUN-2004) to the EMBL/GenBank/DDBJ databases.

## Images



Western blot detection of Ku70 in Hela,A549,MCF7,Jurkat and Raji cell lysates using Ku70 mouse mAb (1:1000 diluted).Predicted band size:70KDa.Observed band size:70KDa.



Immunocytochemistry staining of HeLa cells fixed with 4% Paraformaldehyde and using Ku70 mouse mAb (dilution 1:200).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.