

MYLIP Antibody (Center)

Affinity Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP5315c

Product Information

Application	WB, IHC-P, FC, E
Primary Accession	Q8WY64
Other Accession	NP_037394
Reactivity	Human, Rat, Mouse
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Clone Names	RB18661
Calculated MW	49910
Antigen Region	111-139

Additional Information

Gene ID	29116
Other Names	E3 ubiquitin-protein ligase MYLIP, 632-, Inducible degrader of the LDL-receptor, Idol, Myosin regulatory light chain interacting protein, MIR, MYLIP, BZF1, IDOL
Target/Specificity	This MYLIP antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 111-139 amino acids from the Central region of human MYLIP.
Dilution	WB~~1:1000 IHC-P~~1:100~500 FC~~1:10~50 E~~Use at an assay dependent concentration.
Format	Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.
Storage	Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.
Precautions	MYLIP Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

Protein Information

Name	MYLIP
Synonyms	BZF1, IDOL

Function E3 ubiquitin-protein ligase that mediates ubiquitination and subsequent proteasomal degradation of myosin regulatory light chain (MRLC), LDLR, VLDLR and LRP8. Activity depends on E2 enzymes of the UBE2D family. Proteasomal degradation of MRLC leads to inhibit neurite outgrowth in presence of NGF by counteracting the stabilization of MRLC by saposin-like protein (CNPY2/MSAP) and reducing CNPY2-stimulated neurite outgrowth. Acts as a sterol-dependent inhibitor of cellular cholesterol uptake by mediating ubiquitination and subsequent degradation of LDLR.

Cellular Location Cytoplasm. Cell membrane; Peripheral membrane protein

Tissue Location Ubiquitously expressed.


Background

The ERM protein family members ezrin, radixin, and moesin are cytoskeletal effector proteins linking actin to membrane-bound proteins at the cell surface. Myosin regulatory light chain interacting protein (MYLIP) is a novel ERM-like protein that interacts with myosin regulatory light chain and inhibits neurite outgrowth.


References

Chasman, D.I., et al. PLoS Genet. 5 (11), E1000730 (2009)
 Lindholm, D., et al. Cell. Mol. Life Sci. 66(21):3399-3402(2009)
 Zelcer, N., et al. Science 325(5936):100-104(2009)

Images




All lanes : Anti-MYLIP Antibody (Center) at 1:2000 dilution
 Lane 1: 293 whole cell lysate Lane 2: MCF-7 whole cell lysate
 Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 50 kDa Blocking/Dilution buffer: 5% NFDM/TBST.



MYLIP antibody (Center) (Cat. #AP5315c) immunohistochemistry analysis in formalin fixed and paraffin embedded human brain tissue followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of the MYLIP antibody (Center) for immunohistochemistry. Clinical relevance has not been evaluated.

MYLIP Antibody (Center) (Cat. #AP5315c) flow cytometric analysis of MCF-7 cells (right histogram) compared to a negative control cell (left histogram). FITC-conjugated goat-anti-rabbit secondary antibodies were used for the



analysis.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.