

KDM1/LSD1 Antibody

Purified Mouse Monoclonal Antibody (Mab)
Catalog # AP53268

Product Information

Application	WB, ICC, IP
Primary Accession	O60341
Reactivity	Human
Host	Mouse
Clonality	Monoclonal
Isotype	IgG1
Calculated MW	92903

Additional Information

Gene ID	23028
Other Names	Amine oxidase (flavin containing) domain 2;AOF2;BHC110;BRAF35 HDAC complex protein BHC110;BRAF35-HDAC complex protein BHC110;FAD binding protein BRAF35 HDAC complex, 110 kDa subunit;Flavin-containing amine oxidase domain-containing protein 2;KDM 1;KDM1;Kdm1a;KDM1A_HUMAN;LSD 1;LSD1;Lysine (K) specific demethylase 1;Lysine (K) specific demethylase 1A;Lysine specific histone demethylase 1;Lysine specific histone demethylase 1A;Lysine-specific histone demethylase 1A.
Dilution	WB~~1:1000 ICC~~1:100 IP~~1:500
Format	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide, pH 7.3.
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.

Protein Information

Name	KDM1A (HGNC:29079)
Function	Histone demethylase that can demethylate both 'Lys-4' (H3K4me) and 'Lys-9' (H3K9me) of histone H3, thereby acting as a coactivator or a corepressor, depending on the context (PubMed: 15620353 , PubMed: 15811342 , PubMed: 16079794 , PubMed: 16079795 , PubMed: 16140033 , PubMed: 16223729 , PubMed: 27292636). Acts by oxidizing the substrate by FAD to generate the corresponding imine that is subsequently hydrolyzed (PubMed: 15620353 , PubMed: 15811342 , PubMed: 16079794 , PubMed: 21300290 , PubMed: 26214369). Acts as a corepressor by mediating demethylation of H3K4me, a specific tag for epigenetic transcriptional

activation. Demethylates both mono- (H3K4me1) and di-methylated (H3K4me2) (PubMed:[15620353](#), PubMed:[20389281](#), PubMed:[21300290](#), PubMed:[23721412](#)). May play a role in the repression of neuronal genes. Alone, it is unable to demethylate H3K4me on nucleosomes and requires the presence of RCOR1/CoREST to achieve such activity (PubMed:[16079794](#), PubMed:[16140033](#), PubMed:[16885027](#), PubMed:[21300290](#), PubMed:[23721412](#)). Also acts as a coactivator of androgen receptor (AR)-dependent transcription, by being recruited to AR target genes and mediating demethylation of H3K9me, a specific tag for epigenetic transcriptional repression. The presence of PRKCB in AR-containing complexes, which mediates phosphorylation of 'Thr-6' of histone H3 (H3T6ph), a specific tag that prevents demethylation H3K4me, prevents H3K4me demethylase activity of KDM1A (PubMed:[16079795](#)). Demethylates di-methylated 'Lys-370' of p53/TP53 which prevents interaction of p53/TP53 with TP53BP1 and represses p53/TP53-mediated transcriptional activation. Demethylates and stabilizes the DNA methylase DNMT1 (PubMed:[29691401](#)). Demethylates methylated 'Lys-42' and methylated 'Lys-117' of SOX2 (PubMed:[29358331](#)). Required for gastrulation during embryogenesis. Component of a RCOR/GFI/KDM1A/HDAC complex that suppresses, via histone deacetylase (HDAC) recruitment, a number of genes implicated in multilineage blood cell development (PubMed:[16079794](#), PubMed:[16140033](#)). Facilitates epithelial-to-mesenchymal transition by acting as an effector of SNAI1- mediated transcription repression of epithelial markers E-cadherin/CDH1, CDN7 and KRT8 (PubMed:[20562920](#), PubMed:[27292636](#)). Required for the maintenance of the silenced state of the SNAI1 target genes E-cadherin/CDH1 and CDN7 (PubMed:[20389281](#)). Required for the repression of GIPR expression (PubMed:[34655521](#), PubMed:[34906447](#)).

Cellular Location

Nucleus. Chromosome. Note=Associates with chromatin

Tissue Location

Ubiquitously expressed. [Isoform 4]: Expressed exclusively in brain tissues.

Background


Histone demethylase that demethylates both 'Lys-4' (H3K4me) and 'Lys-9' (H3K9me) of histone H3, thereby acting as a coactivator or a corepressor, depending on the context. Acts by oxidizing the substrate by FAD to generate the corresponding imine that is subsequently hydrolyzed. Acts as a corepressor by mediating demethylation of H3K4me, a specific tag for epigenetic transcriptional activation. Demethylates both mono- (H3K4me1) and di-methylated (H3K4me2) H3K4me. May play a role in the repression of neuronal genes. Alone, it is unable to demethylate H3K4me on nucleosomes and requires the presence of RCOR1/CoREST to achieve such activity. Also acts as a coactivator of androgen receptor (ANDR)-dependent transcription, by being recruited to ANDR target genes and mediating demethylation of H3K9me, a specific tag for epigenetic transcriptional repression. The presence of PRKCB in ANDR-containing complexes, which mediates phosphorylation of 'Thr- 6' of histone H3 (H3T6ph), a specific tag that prevents demethylation H3K4me, prevents H3K4me demethylase activity of KDM1A. Demethylates di-methylated 'Lys-370' of p53/TP53 which prevents interaction of p53/TP53 with TP53BP1 and represses p53/TP53-mediated transcriptional activation. Demethylates and stabilizes the DNA methylase DNMT1. Required for gastrulation during embryogenesis. Component of a RCOR/GFI/KDM1A/HDAC complex that suppresses, via histone deacetylase (HDAC) recruitment, a number of genes implicated in multilineage blood cell development. Effector of SNAI1-mediated transcription repression of E- cadherin/CDH1, CDN7 and KRT8. Required for the maintenance of the silenced state of the SNAI1 target genes E-cadherin/CDH1 and CDN7.

References


- Nagase T.,et al.DNA Res. 5:31-39(1998).
Gregory S.G.,et al.Nature 441:315-321(2006).
Bechtel S.,et al.BMC Genomics 8:399-399(2007).

Hakimi M.-A., et al. Proc. Natl. Acad. Sci. U.S.A. 99:7420-7425(2002).
Humphrey G.W., et al. J. Biol. Chem. 276:6817-6824(2001).


Images



Western blot detection of KDM1/LSD1 in HeLa, HCT116 and COS7 cell lysates using KDM1/LSD1 mouse mAb (1:1000 diluted). Predicted band size: 110 kDa. Observed band size: 110 kDa.



Immunocytochemistry staining of HeLa cells fixed with 4% Paraformaldehyde and using anti-KDM1/LSD1 mouse mAb (dilution 1:100).



Immunoprecipitation analysis of HeLa cell lysates using KDM1/LSD1 mouse mAb.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.