

JMJD6(N-terminus) Antibody

Purified Mouse Monoclonal Antibody (Mab)

Catalog # AP53269

Product Information

Application	WB, ICC
Primary Accession	Q6NYC1
Reactivity	Human
Host	Mouse
Clonality	Monoclonal
Isotype	IgG1
Calculated MW	46462

Additional Information

Gene ID	23210
Other Names	Apoptotic cell clearance receptor;Bifunctional arginine demethylase and lysyl-hydroxylase JMJD6;Histone arginine demethylase JMJD6;JmjC domain-containing protein 6;JMJD 6;JMJD6;JMJD6_HUMAN;Jumonji domain containing 6;jumonji domain-containing protein 6;KIAA0585;Lysyl-hydroxylase JMJD6;Peptide-lysine 5-dioxygenase JMJD6;Phosphatidylserine receptor;Protein PTDSR;PSR;PTDSR 1;PTDSR;PTDSR1.
Dilution	WB~~1:1000 ICC~~1:200
Format	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide, pH 7.3.
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.

Protein Information

Name	JMJD6 (HGNC:19355)
Function	Dioxygenase that can both act as a arginine demethylase and a lysyl-hydroxylase (PubMed: 17947579 , PubMed: 20684070 , PubMed: 21060799 , PubMed: 22189873 , PubMed: 24498420). Acts as a lysyl-hydroxylase that catalyzes 5-hydroxylation on specific lysine residues of target proteins such as U2AF2/U2AF65 and LUC7L2. Regulates RNA splicing by mediating 5-hydroxylation of U2AF2/U2AF65, affecting the pre-mRNA splicing activity of U2AF2/U2AF65 (PubMed: 19574390). Hydroxylates its own N-terminus, which is required for homooligomerization (PubMed: 22189873). Plays a role in the regulation of nucleolar liquid- liquid phase separation (LLPS) by post-translationally modifying LIAT1 at its lysine-rich domain which inhibits LIAT1 nucleolar targeting (By similarity). In addition to peptidyl-lysine

5-dioxygenase activity, may act as an RNA hydroxylase, as suggested by its ability to bind single strand RNA (PubMed:[20679243](#), PubMed:[29176719](#)). Also acts as an arginine demethylase which preferentially demethylates asymmetric dimethylation (PubMed:[17947579](#), PubMed:[24360279](#), PubMed:[24498420](#)). Demethylates histone H3 at 'Arg-2' (H3R2me) and histone H4 at 'Arg-3' (H4R3me), including mono-, symmetric di- and asymmetric dimethylated forms, thereby playing a role in histone code (PubMed:[17947579](#), PubMed:[24360279](#)). However, histone arginine demethylation may not constitute the primary activity in vivo (PubMed:[17947579](#), PubMed:[21060799](#), PubMed:[22189873](#)). In collaboration with BRD4, interacts with the positive transcription elongation factor b (P-TEFb) complex in its active form to regulate polymerase II promoter-proximal pause release for transcriptional activation of a large cohort of genes. On distal enhancers, so called anti-pause enhancers, demethylates both histone H4R3me2 and the methyl cap of 7SKsnRNA leading to the dismissal of the 7SKsnRNA:HEXIM1 inhibitor complex. After removal of repressive marks, the complex BRD4:JMJD6 attract and retain the P-TEFb complex on chromatin, leading to its activation, promoter-proximal polymerase II pause release, and transcriptional activation (PubMed:[24360279](#)). Demethylates other arginine methylated- proteins such as ESR1 (PubMed:[24498420](#)). Has no histone lysine demethylase activity (PubMed:[21060799](#)). Required for differentiation of multiple organs during embryogenesis. Acts as a key regulator of hematopoietic differentiation: required for angiogenic sprouting by regulating the pre-mRNA splicing activity of U2AF2/U2AF65 (By similarity). Seems to be necessary for the regulation of macrophage cytokine responses (PubMed:[15622002](#)).

Cellular Location

Nucleus, nucleoplasm. Nucleus, nucleolus. Cytoplasm. Note=Mainly found throughout the nucleoplasm outside of regions containing heterochromatic DNA, with some localization in nucleolus. During mitosis, excluded from the nucleus and reappears in the telophase of the cell cycle.

Tissue Location

Highly expressed in the heart, skeletal muscle and kidney. Expressed at moderate or low level in brain, placenta, lung, liver, pancreas, spleen, thymus, prostate, testis and ovary. Up- regulated in many patients with chronic pancreatitis. Expressed in nursing thymic epithelial cells.


Background

Dioxygenase that can both act as a histone arginine demethylase and a lysyl-hydroxylase. Acts as a lysyl-hydroxylase that catalyzes 5-hydroxylation on specific lysine residues of target proteins such as U2AF2/U2AF65 and LUC7L2. Acts as a regulator of RNA splicing by mediating 5-hydroxylation of U2AF2/U2AF65, affecting the pre-mRNA splicing activity of U2AF2/U2AF65. In addition to peptidyl-lysine 5-dioxygenase activity, may act as an RNA hydroxylase, as suggested by its ability to bind single strand RNA. Also acts as an arginine demethylase which demethylates histone H3 at 'Arg-2' (H3R2me) and histone H4 at 'Arg-3' (H4R3me), thereby playing a role in histone code. However, histone arginine demethylation may not constitute the primary activity in vivo. Has no histone lysine demethylase activity. Required for differentiation of multiple organs during embryogenesis. Acts as a key regulator of hematopoietic differentiation: required for angiogenic sprouting by regulating the pre-mRNA splicing activity of U2AF2/U2AF65. Seems to be necessary for the regulation of macrophage cytokine responses.


References

- Izawa M.,et al.Submitted (OCT-2001) to the EMBL/GenBank/DDBJ databases.
 Nagase T.,et al.DNA Res. 5:31-39(1998).
 Ota T.,et al.Nat. Genet. 36:40-45(2004).
 Zody M.C.,et al.Nature 440:1045-1049(2006).
 Mural R.J.,et al.Submitted (JUL-2005) to the EMBL/GenBank/DDBJ databases.

Images



Western blot detection of JMJD6(N-terminus) in HeLa and HL-60 lysates using JMJD6(N-terminus) mouse mAb (1:1000 diluted). Predicted band size: 46KDa. Observed band size: 62KDa.



Immunocytochemistry of HeLa cells using anti-JMJD6(N-terminus) mouse mAb diluted 1:200.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.