

beta Actin Antibody

Purified Mouse Monoclonal Antibody (Mab)

Catalog # AP53271

Product Information

| | |
|-------------------|---|
| Application | WB |
| Primary Accession | P60709 |
| Reactivity | Human, Mouse, Rabbit, Neisseria Gonorrhoeae |
| Host | Mouse |
| Clonality | Monoclonal |
| Isotype | IgG2b |
| Calculated MW | 41737 |

Additional Information

| | |
|-------------|--|
| Gene ID | 60 |
| Other Names | A X actin like protein;ACTB;ACTB_HUMAN;Actin;Actin cytoplasmic 1;Actx;b actin;b-actin;Beta cytoskeletal actin;Beta-actin;cytoplasmic 1;Melanoma X actin;N-terminally processed;PS1TP5 binding protein 1;PS1TP5BP1. |
| Dilution | WB~~1:10000 |
| Format | Purified mouse monoclonal in PB(pH 7.4) containing with 0.09% (W/V) sodium azide and 50% glycerol. |
| Storage | Store at -20 °C.Stable for 12 months from date of receipt |

Protein Information

| | |
|----------|---|
| Name | ACTB |
| Function | <p>Actin is a highly conserved protein that polymerizes to produce filaments that form cross-linked networks in the cytoplasm of cells (PubMed:25255767, PubMed:29581253). Actin exists in both monomeric (G-actin) and polymeric (F-actin) forms, both forms playing key functions, such as cell motility and contraction (PubMed:29581253). In addition to their role in the cytoplasmic cytoskeleton, G- and F- actin also localize in the nucleus, and regulate gene transcription and motility and repair of damaged DNA (PubMed:29925947). Plays a role in the assembly of the gamma-tubulin ring complex (gTuRC), which regulates the minus-end nucleation of alpha-beta tubulin heterodimers that grow into microtubule protofilaments (PubMed:39321809, PubMed:38609661). Part of the ACTR1A/ACTB filament around which the dynactin complex is built (By similarity). The dynactin multiprotein complex activates the molecular motor dynein for ultra-processive transport along microtubules (By similarity).</p> |

Cellular Location

Cytoplasm, cytoskeleton. Nucleus Note=Localized in cytoplasmic mRNP granules containing untranslated mRNAs.


Background

Actins are highly conserved proteins that are involved in various types of cell motility and are ubiquitously expressed in all eukaryotic cells.

References

Ponte P.,et al.Nucleic Acids Res. 12:1687-1696(1984).
Nakajima-Iijima S.,et al.Proc. Natl. Acad. Sci. U.S.A. 82:6133-6137(1985).
Ohmori H.,et al.J. Cell Biol. 116:933-941(1992).
Hillier L.W.,et al.Nature 424:157-164(2003).
Gevaert K.,et al.Nat. Biotechnol. 21:566-569(2003).

Images



Western blot detection of beta actin in C6,3T3,COS7,CHO-k1,K562 and Goat muscle cell lysates using beta actin mouse mAb (1:10000 diluted).Predicted band size:45KDa.Observed band size:45KDa.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.