

Anti-CEP78 Antibody

Catalog # AP53295

Product Information

Application	WB, IHC
Primary Accession	Q5JTW2
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	76396

Additional Information

Gene ID	84131
Other Names	C9orf81; Centrosomal protein of 78 kDa; Cep78
Target/Specificity	KLH-conjugated synthetic peptide encompassing a sequence within the center region of human CEP78. The exact sequence is proprietary.
Dilution	WB~~ 1:1000 IHC~~1:100~500
Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.
Storage	Store at -20 °C.Stable for 12 months from date of receipt

Protein Information

Name	CEP78 {ECO:0000303 PubMed:27588451, ECO:0000312 HGNC:HGNC:25740}
Function	Centriole wall protein that localizes to mature centrioles and regulates centriole and cilia biogenesis (PubMed: 27246242 , PubMed: 27588451 , PubMed: 28242748 , PubMed: 34259627). Involved in centrosome duplication: required for efficient PLK4 centrosomal localization and PLK4-induced overduplication of centrioles (PubMed: 27246242). Involved in cilium biogenesis and controls cilium length (PubMed: 27588451). Acts as a regulator of protein stability by preventing ubiquitination of centrosomal proteins, such as CCP110 and tektins (PubMed: 28242748 , PubMed: 34259627). Associates with the EDVP complex, preventing ubiquitination and degradation of CCP110 (PubMed: 28242748 , PubMed: 34259627). Promotes deubiquitination of tektin proteins (TEKT1, TEKT2, TEK3, TEKT4 and TEKT5) via its interaction with USP16 (By similarity).
Cellular Location	Cytoplasm, cytoskeleton, microtubule organizing center, centrosome. Cytoplasm, cytoskeleton, microtubule organizing center, centrosome,

centriole. Cytoplasm, cytoskeleton, cilium basal body Note=Mainly localizes at the centriolar wall, but also found in the pericentriolar material (PubMed:27246242). Expressed in photoreceptor inner segment (PubMed:27588452).


Tissue Location

Widely expressed (PubMed:27588451, PubMed:27588452). Expressed in different retinal cell types with higher expression in cone compared to rod cells (at protein level) (PubMed:27588452).


References

Humphray S.J.,et al.Nature 429:369-374(2004).
Ota T.,et al.Nat. Genet. 36:40-45(2004).
Andersen J.S.,et al.Nature 426:570-574(2003).
Gauci S.,et al.Anal. Chem. 81:4493-4501(2009).
Van Damme P.,et al.Proc. Natl. Acad. Sci. U.S.A. 109:12449-12454(2012).

Images



Western blot analysis of CEP78 expression in COS7 (A), HEK293T (B), CT26 (C), H9C2 (D) whole cell lysates. (Predicted band size: 76 kD; Observed band size: 78 kD)



Immunohistochemical analysis of CEP78 staining in human lung cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.