

MOB3C Antibody

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP53305

Product Information

Application	WB
Primary Accession	Q70IA8
Reactivity	Human, Mouse
Host	Rabbit
Clonality	Polyclonal
Calculated MW	25623

Additional Information

Gene ID	148932
Other Names	MOB kinase activator 3C, Mob1 homolog 2C, Mps one binder kinase activator-like 2C, MOB3C, MOBKL2C
Target/Specificity	KLH-conjugated synthetic peptide encompassing a sequence within the center region of human MOB3C. The exact sequence is proprietary.
Dilution	WB~~ 1:1000
Format	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.09% (W/V) sodium azide and 50% glycerol
Storage	Store at -20 °C.Stable for 12 months from date of receipt

Protein Information

Name	MOB3C
Synonyms	MOBKL2C
Function	May regulate the activity of kinases.

Background


May regulate the activity of kinases.

References

Florindo C.S.,et al.Submitted (AUG-2003) to the EMBL/GenBank/DDBJ databases.

Jikuya H.,et al.Submitted (JUL-2002) to the EMBL/GenBank/DDBJ databases.
Gregory S.G.,et al.Nature 441:315-321(2006).
Mural R.J.,et al.Submitted (SEP-2005) to the EMBL/GenBank/DDBJ databases.
Ota T.,et al.Nat. Genet. 36:40-45(2004).

Images



Anti-MOB3C Antibody at 1:1000 dilution + K562 whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 26 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.