

# SAR1B Antibody

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP53321

## Product Information

Application	WB
Primary Accession	<a href="#">Q9Y6B6</a>
Reactivity	Human, Mouse
Host	Rabbit
Clonality	Polyclonal
Calculated MW	22410

## Additional Information

Gene ID	51128
Other Names	GTP-binding protein SAR1b, GTP-binding protein B, GTBPB, SAR1B, SARA2, SARB
Target/Specificity	KLH-conjugated synthetic peptide encompassing a sequence within the center region of human SAR1B. The exact sequence is proprietary.
Dilution	WB~~ 1:1000
Format	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.09% (W/V) sodium azide and 50% glycerol
Storage	Store at -20 °C.Stable for 12 months from date of receipt

## Protein Information

Name	SAR1B {ECO:0000303   PubMed:33186557, ECO:0000312   HGNC:HGNC:10535}
Function	<p>Small GTPase that cycles between an active GTP-bound and an inactive GDP-bound state and mainly functions in vesicle-mediated endoplasmic reticulum (ER) to Golgi transport. The active GTP-bound form inserts into the endoplasmic reticulum membrane where it recruits the remainder of the coat protein complex II/COPII (PubMed:<a href="#">23433038</a>, PubMed:<a href="#">32358066</a>, PubMed:<a href="#">33186557</a>, PubMed:<a href="#">36369712</a>). The coat protein complex II assembling and polymerizing on endoplasmic reticulum membrane is responsible for both the sorting of cargos and the deformation and budding of membranes into vesicles destined to the Golgi (PubMed:<a href="#">23433038</a>, PubMed:<a href="#">32358066</a>, PubMed:<a href="#">33186557</a>). In contrast to SAR1A, SAR1B specifically interacts with the cargo receptor SURF4 to mediate the transport of lipid-carrying lipoproteins including APOB and APOA1 from the endoplasmic reticulum to the Golgi and thereby, indirectly regulates lipid</p>

homeostasis (PubMed:[32358066](#), PubMed:[33186557](#)). In addition to its role in vesicle trafficking, can also function as a leucine sensor regulating TORC1 signaling and more indirectly cellular metabolism, growth and survival. In absence of leucine, interacts with the GATOR2 complex via MIOS and inhibits TORC1 signaling. The binding of leucine abrogates the interaction with GATOR2 and the inhibition of the TORC1 signaling. This function is completely independent of the GTPase activity of SAR1B (PubMed:[34290409](#)).

#### Cellular Location

Endoplasmic reticulum membrane; Peripheral membrane protein {ECO:0000250|UniProtKB:Q9QVY3}. Golgi apparatus, Golgi stack membrane {ECO:0000250|UniProtKB:Q9QVY3}; Peripheral membrane protein {ECO:0000250|UniProtKB:Q9QVY3}. Cytoplasm, cytosol. Lysosome membrane. Note=Active at endoplasmic reticulum exit sites (ERES) where it inserts into the membrane and recruits the remainder of the coat protein complex II/COPII (PubMed:23433038). Upon leucine deprivation, associates with lysosomal membranes to repress TORC1 signaling (PubMed:34290409).

#### Tissue Location

Expressed in many tissues including small intestine, liver, muscle and brain.


## Background

Involved in transport from the endoplasmic reticulum to the Golgi apparatus. Activated by the guanine nucleotide exchange factor PREB. Involved in the selection of the protein cargo and the assembly of the COPII coat complex.

## References

Song H.,et al.Submitted (SEP-1998) to the EMBL/GenBank/DDBJ databases.  
Zhou Y.,et al.Submitted (JUL-2003) to the EMBL/GenBank/DDBJ databases.  
Mural R.J.,et al.Submitted (SEP-2005) to the EMBL/GenBank/DDBJ databases.  
Burkard T.R.,et al.BMC Syst. Biol. 5:17-17(2011).  
Jones B.,et al.Nat. Genet. 34:29-31(2003).

## Images



Anti-SAR1B Antibody at 1:1000 dilution + HeLa whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 22 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.