

# CstF-64T Antibody

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP53338

## Product Information

Application	WB
Primary Accession	<a href="#">Q9H0L4</a>
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	64437

## Additional Information

Gene ID	23283
Other Names	Cleavage stimulation factor subunit 2 tau variant, CF-1 64 kDa subunit tau variant, Cleavage stimulation factor 64 kDa subunit tau variant, CSTF 64 kDa subunit tau variant, TauCstF-64, CSTF2T, KIAA0689
Dilution	WB~~ 1:1000
Format	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.09% (W/V) sodium azide and 50% glycerol
Storage	Store at -20 °C.Stable for 12 months from date of receipt

## Protein Information

Name	CSTF2T
Synonyms	KIAA0689
Function	May play a significant role in AAUAAA-independent mRNA polyadenylation in germ cells. Directly involved in the binding to pre- mRNAs (By similarity).
Cellular Location	Nucleus.

## Background


May play a significant role in AAUAAA-independent mRNA polyadenylation in germ cells. Directly involved in the binding to pre-mRNAs (By similarity).

## References

Dass B.,et al.Genomics 80:509-514(2002).  
Wiemann S.,et al.Genome Res. 11:422-435(2001).  
Ota T.,et al.Nat. Genet. 36:40-45(2004).  
Suzuki Y.,et al.Submitted (APR-2005) to the EMBL/GenBank/DDBJ databases.  
Mural R.J.,et al.Submitted (JUL-2005) to the EMBL/GenBank/DDBJ databases.

## Images

---



Anti-CstF-64T Antibody at 1:1000 dilution + Jurkat whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 64 kDa  
Blocking/Dilution buffer: 5% NFDM/TBST.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.