

# Cytochrome P450 2W1 Antibody

Purified Rabbit Polyclonal Antibody (Pab)

Catalog # AP53339

## Product Information

Application	WB
Primary Accession	<a href="#">Q8TAV3</a>
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	53844

## Additional Information

Gene ID	54905
Other Names	Cytochrome P450 2W1, 1.14.14.-, CYP2W1
Dilution	WB~~ 1:1000
Format	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.09% (W/V) sodium azide and 50% glycerol
Storage	Store at -20 °C.Stable for 12 months from date of receipt

## Protein Information

Name	CYP2W1 {ECO:0000303 PubMed:26936974, ECO:0000312 HGNC:HGNC:20243}
Function	<p>A cytochrome P450 monooxygenase that may play a role in retinoid and phospholipid metabolism (PubMed:<a href="#">22591743</a>, PubMed:<a href="#">26936974</a>). Catalyzes the hydroxylation of saturated carbon hydrogen bonds. Hydroxylates all trans-retinoic acid (atRA) to 4- hydroxyretinoate and may regulate atRA clearance. Other retinoids such as all-trans retinol and all-trans retinal are potential endogenous substrates (PubMed:<a href="#">26936974</a>). Catalyzes both epoxidation of double bonds and hydroxylation of carbon hydrogen bonds of the fatty acyl chain of 1-acylphospholipids/2-lysophospholipids. Can metabolize various lysophospholipids classes including lysophosphatidylcholines (LPCs), lysophosphatidylinositols (LPIs), lysophosphatidylserines (LPSs), lysophosphatidylglycerols (LPGs), lysophosphatidylethanolamines (LPEs) and lysophosphatidic acids (LPAs) (PubMed:<a href="#">22591743</a>). Has low or no activity toward 2-acylphospholipids/1-lysophospholipids, diacylphospholipids and free fatty acids (PubMed:<a href="#">22591743</a>, PubMed:<a href="#">26936974</a>). May play a role in tumorigenesis by activating procarcinogens such as aflatoxin B1, polycyclic aromatic hydrocarbon dihydrodiols and aromatic amines (PubMed:<a href="#">16551781</a>,</p>

PubMed:[20805301](#), PubMed:[24278521](#)). Mechanistically, uses molecular oxygen inserting one oxygen atom into a substrate, and reducing the second into a water molecule, with two electrons provided by NADPH via cytochrome P450 reductase (CPR; NADPH-ferrihemoprotein reductase) (PubMed:[22591743](#), PubMed:[26936974](#)).

**Cellular Location**

Endoplasmic reticulum lumen. Cell membrane. Microsome membrane.  
Note=About 8% are expressed on the cell surface.

**Tissue Location**

Very low levels are detected in fetal and adult tissues. Highly expressed in several tumor samples, in particular colon and adrenal tumors.


## Background

Seems to have broad catalytic activity towards several chemicals, including polycyclic aromatic hydrocarbon dihydrodiols and aromatic amines (PubMed:[16551781](#), PubMed:[24278521](#)). Active also in the metabolism of indoline substrates and is able to activate aflatoxin B1 into cytotoxic products (PubMed:[20805301](#)). Furthermore, it seems to be involved in the oxidation of lysophospholipids and fatty acids (PubMed:[22591743](#)).

## References

Hillier L.W., et al. *Nature* 424:157-164(2003).  
Karlgrén M., et al. *Biochem. Biophys. Res. Commun.* 341:451-458(2006).  
Wu Z.L., et al. *Mol. Pharmacol.* 69:2007-2014(2006).  
Gomez A., et al. *Mol. Pharmacol.* 78:1004-1011(2010).  
Eun C.Y., et al. *Toxicol. Res.* 26:171-175(2010).

## Images



Anti-Cytochrome P450 2W1 Antibody at 1:1000 dilution + HepG2 whole cell lysate. Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 54 kDa. Blocking/Dilution buffer: 5% NFDM/TBST.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.