10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

Anti-Renin Antibody

Catalog # AP53367

Product Information

Application	WB, IHC, IF/IC
Primary Accession	P00797
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	45057

Additional Information

Gene ID	5972
Other Names	Renin; Angiotensinogenase
Target/Specificity	KLH-conjugated synthetic peptide encompassing a sequence within the center region of human Renin. The exact sequence is proprietary.
Dilution	WB~~ 1:1000 IHC~~1:100~500 IF/IC~~N/A
Format	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.
Storage	Store at -20 °C.Stable for 12 months from date of receipt

Protein Information


Name	REN
Function	Renin is a highly specific endopeptidase, whose only known function is to generate angiotensin I from angiotensinogen in the plasma, initiating a cascade of reactions that produce an elevation of blood pressure and increased sodium retention by the kidney.
Cellular Location	Secreted. Membrane. Note=Associated to membranes via binding to ATP6AP2.

References


Imai T.,et al.Proc. Natl. Acad. Sci. U.S.A. 80:7405-7409(1983).
Morris B.J.,et al.Clin. Sci. 71:345-355(1986).
Hardman J.A.,et al.DNA 3:457-468(1984).
Rieder M.J.,et al.Submitted (OCT-2003) to the EMBL/GenBank/DDBJ databases.

Ebert L.,et al.Submitted (JUN-2004) to the EMBL/GenBank/DDBJ databases.


Images



Western blot analysis of Renin expression in K562 (A), Jurkat (B), HEK293T (C) whole cell lysates. (Predicted band size: 45 kD; Observed band size: 38 kD)



Immunohistochemical analysis of Renin staining in human placenta formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



Immunofluorescent analysis of Renin staining in MCF7 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a hidified chamber. Cells were washed with PBST and incubated with a AF488-conjugated secondary antibody (green) in PBS at room temperature in the dark. Phalloidin - AF594 was used to stain Actin filaments (red). DAPI was used to stain the cell nuclei (blue).

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.