10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

# Anti-BiP/GRP78 Antibody

Mouse Anti Human Monoclonal Antibody
Catalog # AP53389

## Product Information

| | |
|---|---|
| **Application** | WB, IF, IP |
| **Primary Accession** | P11021 |
| **Other Accession** | NM_005347 |
| **Reactivity** | Human, Mouse, Rat |
| **Host** | Mouse |
| **Clonality** | Monoclonal |
| **Isotype** | IgG2b |
| **Immunogen** | Purified recombinant human BiP/GRP78 protein expressed in E.coli. |
| **Purification** | Affinity purified |
| **Calculated MW** | 72333 |

## Additional Information

| | |
|---|---|
| **Gene ID** | 3309 |
| **Other Names** | 78 kDa glucose regulated protein;78 kDa glucose-regulated protein;AL022860;AU019543;BIP;D2Wsu141e;D2Wsu17e;Endoplasmic reticulum lumenal Ca(2+)-binding protein grp78;Endoplasmic reticulum lumenal Ca2+ binding protein grp78;FLJ26106;Glucose Regulated Protein 78kDa;GRP 78;GRP-78;GRP78;GRP78_HUMAN;Heat shock 70 kDa protein 5;Heat Shock 70kDa Protein 5;Hsce70;HSPA 5;HSPA5;Immunoglobulin Heavy Chain Binding Protein;Immunoglobulin heavy chain-binding protein;mBiP;MIF2;Sez7. |
| **Dilution** | WB~~1:1000 IF~~1:50~200 IP~~N/A |
| **Format** | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide, pH 7.3. |
| **Storage** | Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles. |

## Protein Information

| | |
|---|---|
| **Name** | HSPA5 ( HGNC:5238) |
| **Function** | Endoplasmic reticulum chaperone that plays a key role in protein folding and quality control in the endoplasmic reticulum lumen (PubMed:2294010, PubMed:23769672, PubMed:23990668, PubMed:28332555). Involved in the correct folding of proteins and degradation of misfolded proteins via its interaction with DNAJC10/ERdj5, probably to facilitate the release of |

DNAJC10/ERdj5 from its substrate (By similarity). Acts as a key repressor of the EIF2AK3/PERK and ERN1/IRE1- mediated unfolded protein response (UPR) (PubMed:11907036, PubMed:1550958, PubMed:19538957, PubMed:36739529). In the unstressed endoplasmic reticulum, recruited by DNAJB9/ERdj4 to the luminal region of ERN1/IRE1, leading to disrupt the dimerization of ERN1/IRE1, thereby inactivating ERN1/IRE1 (By similarity). Also binds and inactivates EIF2AK3/PERK in unstressed cells (PubMed:11907036). Accumulation of misfolded protein in the endoplasmic reticulum causes release of HSPA5/BiP from ERN1/IRE1 and EIF2AK3/PERK, allowing their homodimerization and subsequent activation (PubMed:11907036). Plays an auxiliary role in post-translational transport of small presecretory proteins across endoplasmic reticulum (ER). May function as an allosteric modulator for SEC61 channel-forming translocon complex, likely cooperating with SEC62 to enable the productive insertion of these precursors into SEC61 channel. Appears to specifically regulate translocation of precursors having inhibitory residues in their mature region that weaken channel gating. May also play a role in apoptosis and cell proliferation (PubMed:26045166).


**Cellular Location**

Endoplasmic reticulum lumen. Melanosome. Cytoplasm {ECO:0000250|UniProtKB:P20029}. Cell surface Note=Identified by mass spectrometry in melanosome fractions from stage I to stage IV (PubMed:12643545). Localizes to the cell surface of epithelial cells in response to high levels of free iron (PubMed:20484814, PubMed:24355926, PubMed:27159390)


## Background

Probably plays a role in facilitating the assembly of multimeric protein complexes inside the endoplasmic reticulum. Involved in the correct folding of proteins and degradation of misfolded proteins via its interaction with DNAJC10, probably to facilitate


## Images




Western blot analysis of extracts from 3T3,A549,C6 and Hela cell lysates using BiP/GRP78 mouse mAb (1:1000 diluted).Predicted band size:72KDa.Observed band size:72KDa.



Immunofluorescent analysis of Hela cells fixed fixed by anhydrous methanol at -20℃ and using BiP/GRP78 mouse mAb (dilution 1:200).




Immunoprecipitation of BiP/GRP78 from Hela cell extracts using BiP/GRP78 Mouse mAb.Western blot was performed using BiP/GRP78 Mouse mAb.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.