

Histone H2B (7E2) Mouse mAb

Catalog # AP53518

Product Information

Application	WB, IHC
Primary Accession	P33778
Reactivity	Rat
Host	Mouse
Clonality	Monoclonal Antibody
Calculated MW	13950

Additional Information

Gene ID	3018
Other Names	H2B GL105, H2B.1, Histone H2B type 1B, HIST1H2BB
Dilution	WB~~1:100000 IHC~~1:100~500

Protein Information


Name	H2BC3 (HGNC:4751)
Function	Core component of nucleosome. Nucleosomes wrap and compact DNA into chromatin, limiting DNA accessibility to the cellular machineries which require DNA as a template. Histones thereby play a central role in transcription regulation, DNA repair, DNA replication and chromosomal stability. DNA accessibility is regulated via a complex set of post-translational modifications of histones, also called histone code, and nucleosome remodeling.
Cellular Location	Nucleus. Chromosome.

Background


Swiss-Prot Acc.P33778.Core component of nucleosome. Nucleosomes wrap and compact DNA into chromatin, limiting DNA accessibility to the cellular machineries which require DNA as a template. Histones thereby play a central role in transcription regulation, DNA repair, DNA replication and chromosomal stability. DNA accessibility is regulated via a complex set of post-translational modifications of histones, also called histone code, and nucleosome remodeling.

Images

Western blot detection of Histone H2B in Hela and COS7 cell lysates using Histone H2B mouse mAb (1:100000)



diluted). Predicted band size: 14 kDa. Observed band size: 14 kDa.



Immunohistochemical analysis of paraffin-embedded human breast carcinoma using Histone H2B mouse mAb diluted at 1:500.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.