

HLTF Rabbit pAb

Catalog # AP53577

Product Information

Primary Accession	Q14527
Host	Rabbit
Clonality	Polyclonal Antibody
Calculated MW	113929

Additional Information

Gene ID	6596
Other Names	ZBU1; HLTF1; RNF80; HIP116; SNF2L3; HIP116A; SMARCA3
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.

Protein Information

Name	HLTF (HGNC:11099)
Function	Functions as a DNA-dependent ATPase and E3 ubiquitin-protein ligase involved in chromatin regulation and DNA damage tolerance (DDT) (PubMed:18316726, PubMed:18719106, PubMed:26051180, PubMed:31960921, PubMed:39142279, PubMed:40680746). Catalyzes 'Lys-63'-linked polyubiquitination of monoubiquitinated PCNA at 'Lys-164' in response to genotoxic stress, promoting error-free postreplication repair via template switching (PubMed:18316726, PubMed:18719106). Acts as an epigenetic regulator by promoting recruitment of DNMT1, thereby ensuring DNA methylation inheritance: specifically binds histone H3 trimethylated at 'Lys-9' (H3K9me3) and mediates histone H3 'Lys-23' polyubiquitination (H3K23ub), a docking site for DNMT1, leading to DNMT1 recruitment and replication-coupled DNA methylation maintenance (PubMed:40680746). Catalyzes formation of H3K23ub in two steps: first mediates monoubiquitination together with UBE2E1 and UBE2D2, and then extends ubiquitin chains via 'Lys-63'-linked ubiquitination together with UBE2N and UBE2V2 (PubMed:40680746). Also acts as a chromatin redodeling factor, thereby regulating transcription (PubMed:10391891, PubMed:1994885, PubMed:9126292). Exhibits ATP-dependent double-stranded DNA (dsDNA) translocase activity but lacks classical helicase activity; mediates replication fork reversal by concertedly unwinding and annealing nascent and parental strands, thereby suppressing DNA synthesis and maintaining genomic stability (PubMed:1994885). Resolves G-quadruplex (G4) DNA structures in cooperation with MSH2, limiting replication stress and G4 accumulation across the cell cycle (PubMed:39142279). Contributes to nucleotide excision repair by evicting lesion-containing oligonucleotides using its HIRAN and

ATPase domains (PubMed:[26051180](#)). Can displace single-stranded DNA from triplex structures through ATP-dependent dsDNA translocation (PubMed:[26051180](#), PubMed:[31960921](#)). Also has protein clearing activity at the stalled replication fork, facilitating restart of DNA replication (PubMed:[21795603](#)).

Cellular Location

Nucleus. Chromosome


Tissue Location

Expressed in brain, heart, kidney, liver, lung, pancreas, placenta and skeletal muscle.

Background

Swiss-Prot Acc.Q14527.Has both helicase and E3 ubiquitin ligase activities. Possesses intrinsic ATP-dependent nucleosome-remodeling activity; This activity may be required for transcriptional activation or repression of specific target promoters . These may include the SERPINE1 and HIV-1 promoters and the SV40 enhancer, to which this protein can bind directly. Plays a role in error-free postreplication repair (PRR) of damaged DNA and maintains genomic stability through acting as a ubiquitin ligase for 'Lys-63'-linked polyubiquitination of chromatin-bound PCNA.

Images



Western blot detection of HLTF in HeLa cell lysates using HLTF Rabbit pAb(1:1000 diluted). Predicted band size:114kDa.Observed band size:114kDa.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.