

HMGCR Rabbit pAb

Catalog # AP53578

Product Information

Primary Accession	P04035
Host	Rabbit
Clonality	Polyclonal Antibody
Calculated MW	97476

Additional Information

Gene ID	3156
Other Names	LDLCQ3
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.


Protein Information

Name	HMGCR (HGNC:5006)
Function	Catalyzes the conversion of (3S)-hydroxy-3-methylglutaryl-CoA (HMG-CoA) to mevalonic acid, the rate-limiting step in the synthesis of cholesterol and other isoprenoids, thus plays a critical role in cellular cholesterol homeostasis (PubMed: 21357570 , PubMed: 2991281 , PubMed: 36745799 , PubMed: 6995544). HMGCR is the main target of statins, a class of cholesterol-lowering drugs (PubMed: 11349148 , PubMed: 18540668 , PubMed: 36745799).
Cellular Location	Endoplasmic reticulum membrane; Multi-pass membrane protein {ECO:0000250 UniProtKB:P00347}. Peroxisome membrane; Multi-pass membrane protein {ECO:0000250 UniProtKB:P00347}
Tissue Location	[Isoform 1]: Ubiquitously expressed with the highest levels in the cerebellum, fetal brain, testis, skin and adrenal gland. [Isoform 3]: Low abundance except in skin, esophagus, and uterine cervix.

Background

Swiss-Prot Acc.P04035.Transmembrane glycoprotein that is the rate-limiting enzyme in cholesterol biosynthesis as well as in the biosynthesis of nonsterol isoprenoids that are essential for normal cell function including ubiquinone and geranylgeranyl proteins.

Images



Western blot detection of HMGCR in K562,3T3,Hela cell lysates using HMGCR Rabbit pAb(1:1000 diluted).Predicted band size:98kDa.Observed band size:98kDa.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.