

XPO4 Rabbit pAb

Catalog # AP53627

Product Information

Primary Accession	Q9C0E2
Host	Rabbit
Clonality	Polyclonal Antibody
Calculated MW	130139

Additional Information

Gene ID	64328
Other Names	exp4
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.


Protein Information

Name	XPO4 {ECO:0000303 PubMed:26787900, ECO:0000312 HGNC:HGNC:17796}
Function	Mediates the nuclear export of proteins (cargos), such as EIF5A, SMAD3 and isoform M2 of PKM (PKM2) (PubMed:10944119, PubMed:16449645, PubMed:26787900). In the nucleus binds cooperatively to its cargo and to the GTPase Ran in its active GTP-bound form. Docking of this trimeric complex to the nuclear pore complex (NPC) is mediated through binding to nucleoporins (PubMed:10944119, PubMed:16449645). Upon transit of a nuclear export complex into the cytoplasm, disassembling of the complex and hydrolysis of Ran-GTP to Ran-GDP (induced by RANBP1 and RANGAP1, respectively) cause release of the cargo from the export receptor (PubMed:10944119, PubMed:16449645). XPO4 then return to the nuclear compartment and mediate another round of transport (PubMed:10944119, PubMed:16449645). The directionality of nuclear export is thought to be conferred by an asymmetric distribution of the GTP- and GDP-bound forms of Ran between the cytoplasm and nucleus (PubMed:10944119, PubMed:16449645). Catalyzes the nuclear export of hypusinated EIF5A; a small cytoplasmic protein that enters nucleus and accumulates within nucleolus if not exported back by XPO4 (PubMed:10944119). Specifically mediates nuclear export of isoform M2 of PKM (PKM2) following PKM2 deacetylation by SIRT6 (PubMed:26787900). Also mediates the nuclear import of SOX transcription factors SRY and SOX2 (By similarity).
Cellular Location	Cytoplasm. Nucleus. Note=Shuttles between the nucleus and the cytoplasm.

Background

Swiss-Prot Acc.Q9C0E2.Mediates the nuclear export of proteins (cargos) with broad substrate specificity. In the nucleus binds cooperatively to its cargo and to the GTPase Ran in its active GTP-bound form. Docking of this trimeric complex to the nuclear pore complex (NPC) is mediated through binding to nucleoporins. Upon transit of a nuclear export complex into the cytoplasm, disassembling of the complex and hydrolysis of Ran-GTP to Ran-GDP (induced by RANBP1 and RANGAP1, respectively) cause release of the cargo from the export receptor. XPO4 then return to the nuclear compartment and mediate another round of transport. The directionality of nuclear export is thought to be conferred by an asymmetric distribution of the GTP- and GDP-bound forms of Ran between the cytoplasm and nucleus.

Images



Western blot detection of Exportin 4 in MCF7,C6 cell lysates using Exportin 4 Rabbit pAb(1:1000 diluted).Predicted band size:130KDa.Observed band size:130KDa.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.