10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

TPPP Rabbit pAb

Catalog # AP53634

Product Information

Primary Accession	O94811
Host	Rabbit
Clonality	Polyclonal Antibody
Calculated MW	23694

Additional Information

Gene ID	11076
Other Names	p24; p25; TPPP1; TPPP/p25; p25alpha
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.

Protein Information

Name	TPPP {ECO:0000303|PubMed:17105200, ECO:0000312|HGNC:HGNC:24164}
Function	Regulator of microtubule dynamics that plays a key role in myelination by promoting elongation of the myelin sheath (PubMed:31522887). Acts as a microtubule nucleation factor in oligodendrocytes: specifically localizes to the postsynaptic Golgi apparatus region, also named Golgi outpost, and promotes microtubule nucleation, an important step for elongation of the myelin sheath (PubMed:31522887, PubMed:33831707). Required for both uniform polarized growth of distal microtubules as well as directing the branching of proximal processes (PubMed:31522887). Shows magnesium-dependent GTPase activity; the role of the GTPase activity is unclear (PubMed:21316364, PubMed:21995432). In addition to microtubule nucleation activity, also involved in microtubule bundling and stabilization of existing microtubules, thereby maintaining the integrity of the microtubule network (PubMed:17105200, PubMed:17693641, PubMed:18028908, PubMed:26289831). Regulates microtubule dynamics by promoting tubulin acetylation: acts by inhibiting the tubulin deacetylase activity of HDAC6 (PubMed:20308065, PubMed:23093407). Also regulates cell migration: phosphorylation by ROCK1 inhibits interaction with HDAC6, resulting in decreased acetylation of tubulin and increased cell motility (PubMed:23093407). Plays a role in cell proliferation by regulating the G1/S-phase transition (PubMed:23355470). Involved in astral microtubule organization and mitotic spindle orientation during early stage of mitosis; this process is regulated by phosphorylation by LIMK2 (PubMed:22328514).
Cellular Location	Golgi outpost {ECO:0000250|UniProtKB:D3ZQL7}. Cytoplasm, cytoskeleton, microtubule organizing center {ECO:0000250|UniProtKB:D3ZQL7}. Cytoplasm,


cytoskeleton. Nucleus Cytoplasm, cytoskeleton, spindle Note=Specifically localizes to the postsynaptic Golgi apparatus region, also named Golgi outpost, which shapes dendrite morphology by functioning as sites of acentrosomal microtubule nucleation (By similarity). Mainly localizes to the cytoskeleton (PubMed:18028908) Also found in the nucleus; however, nuclear localization is unclear and requires additional evidence (PubMed:18028908). Localizes to glial Lewy bodies in the brains of individuals with synucleinopathies (PubMed:15590652, PubMed:17027006). During mitosis, colocalizes with LIMK2 at the mitotic spindle (PubMed:22328514) {ECO:0000250|UniProtKB:D3ZQL7, ECO:0000269|PubMed:15590652, ECO:0000269|PubMed:17027006, ECO:0000269|PubMed:18028908, ECO:0000269|PubMed:22328514}

Tissue Location Widely expressed..

Background

Swiss-Prot Acc.O94811.May play a role in the polymerization of tubulin into microtubules, microtubule bundling and the stabilization of existing microtubules, thus maintaining the integrity of the microtubule network. May play a role in mitotic spindle assembly and nuclear envelope breakdown.

Images



Western blot detection of TPPP in Rat Brain lysates using TPPP Rabbit pAb(1:1000 diluted).Predicted band size:24kDa.Observed band size:24kDa.

Please note: All products are 'FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC OR THERAPEUTIC PROCEDURES'.